



**ANNUAL
REPORT
2018**



ANNUAL REPORT

2018

02 – 03

FOREWORD

04 – 41

RESEARCH
GROUPS

42 – 51

CORE FACILITIES

52 – 75

FACTS &
FIGURES

FOREWORD

OVERVIEW

2018 brought some excellent news for the future of IMB with an additional commitment from the Boehringer Ingelheim Foundation and the State of Rhineland-Palatinate to jointly fund IMB with €106 million until 2027. Three new Group Leaders joined IMB in the last year with one more joining in 2019. During the year, IMB hosted several successful international events including an EMBO workshop. IMB has been heavily involved in preparing an application for a Collaborative Research Center (CRC), which was greenlit by the DFG in November. The developments of 2018 are shaping the future of IMB, consolidating IMB's position as a leading international research centre.

NEW DEVELOPMENTS AT IMB

This year saw two major developments at IMB. First and foremost, IMB's future was consolidated with the announcement from the Boehringer Ingelheim Foundation and the state of Rhineland-Palatinate of the continuation of IMB's core funding beyond its first funding period. Starting in autumn 2020, the foundation will contribute €54 million and the state €52 million. This second round of financing secures IMB's core funding until mid-2027. The funding follows on from the generous donation of €100 million by the Boehringer Ingelheim Foundation in 2010, which provided core funding for IMB through the period 2010 to 2020. With this funding secured, IMB can continue recruiting top talents in the life sciences while providing state of the art facilities and technology to its researchers. In November this year, the DFG announced €12.4 million in funding for a Collaborative Research Center on "DNA Repair and Genome Stability", which will be coordinated by IMB and comprises 8 projects from IMB group leaders, 3 support services provided by IMB, and an Integrated Research Training Group that will link in with the existing International PhD Programme at IMB. The initiative brings together researchers from IMB, Johannes Gutenberg University Mainz, University Medical Center Mainz, University of Technology Darmstadt, Ludwig Maximilian University Munich, and Goethe University Frankfurt to create a research network on DNA repair and genome stability in which IMB will be a central hub.



STAFF CHANGES

2018 saw three new Group Leaders join IMB: Peter Baumann, Edward Lemke and Anton Khmelinskii. Peter joins us from the Stowers Institute, Kansas as an Adjunct Director and an Alexander von Humboldt Professor at JGU. Peter's research focuses on both telomere biology and chromosomal inheritance. Edward is also a new Adjunct Director and Professor of synthetic biophysics at JGU. Edward and his lab will be moving to Mainz from EMBL, Heidelberg upon completion of the second new university biology building (Biocenter II) in late 2019. Edward's group probes the structure and function of intrinsically disordered proteins. Anton joins IMB as a new Junior Group Leader from the Center for Molecular Biology in Heidelberg. Anton investigates protein quality control systems and the contributions of protein ubiquitylation to these. Furthermore, Martin Möckel has been appointed as the new head of our Protein Production Core Facility in March 2018.

We are also delighted to announce that Joan Barau from the Institut Curie in Paris will be joining IMB in 2019. Joan was a recipient of the French Academy of Sciences prize for great advances in biological sciences in 2015. He specialises in developmental epigenetics and genome stability and will fit in extremely well with the current research focus at IMB. A further round of recruitment for two additional group leader positions will commence next year. We also say goodbye to three of our members this year: Miguel Andrade, Holger Richly and Natalia Soshnikova. I wish them all the best with their future plans.



RESEARCH ACTIVITIES AND EVENTS

2018 was one of the busiest but most productive years for IMB to date. Three international events were held at IMB this year. In March, we hosted a one-day symposium on "Gene Regulation in Evolution" which aimed at bridging the gap between researchers focusing on the molecular side of gene regulation and evolutionary biologists. In October, IMB held a four-day EMBO workshop on "RNA and Genome Maintenance: Cooperation and Conflict Management". The workshop featured 25 speakers and 15 short talks, with 187 participants from 30 countries attending. In November, we had a workshop on "Molecular Mechanisms of Circadian Clocks", which featured talks from 16 renowned international speakers. In addition to these events, IMB also hosted 17 seminars from distinguished speakers, including three organised by IMB's postdocs, and several outreach activities.

IMB continued with its trend of publishing excellent research with 81 research publications in 2018. Highlight articles from this year include publications from the Ketting group in *Developmental Cell*, the Butter group in *Science* and the Niehrs group in *Nature Genetics*. Furthermore, Natalia Soshnikova was accepted into the prestigious Heisenberg Programme of the DFG. Christof Niehrs was the recipient of an ERC Advanced Grant to pursue research into epigenetic regulation via R-loops. My own research was supported with the awarding of an ERC Proof of Concept Grant to explore the translation of a ubiquitin toolkit, developed in my group, into a commercial product.

Our community of PhD students and postdocs continues to thrive at IMB. The international PhD Programme now consists of 128 students from 31 different countries and this year celebrated its 40th graduate. As part of the IMB Postdoc Programme, we also launched our first coordinated recruitment call for postdoctoral researchers. With interviews taking place in early 2019, we expect this format to recruit exceptional junior scientists to IMB. The 7th International Summer School at IMB retained its track record of offering great training and hosted 17 international students this summer.


OUTLOOK

For 2019 we have already planned our IMB Conference on "Chromosome Territories and Nuclear Architecture" in October. This event is in honour of Christoph Cremer who will be retiring at the end of next year. The Collaborative Research Center goes into full swing next year and IMB will host the kick-off meeting for the new programme at the end of summer. In March, we will be holding a practical course in the context of a pan-European initiative, UbiCODE. Finally, in May, Christof Niehrs will organise a public event in cooperation with the Academy of Science and Literature, Mainz on the ethics of CRISPR/Cas technology.

I would like to thank the Boehringer Ingelheim Foundation for their support and generous continued funding of our institute. I wish to say a special thank you to the members of our Scientific Advisory Board (SAB) who have been instrumental in the growth and success of IMB. In particular, I wish to sincerely welcome Geneviève Almouzni from the Institut Curie who has joined our SAB this year. Finally, I am very grateful to all the staff at IMB who have made my first year as Executive Director as smooth as possible.

Helle Ulrich Executive Director

RESEARCH GROUPS



ANDRADE
PAGE 06



BAUMANN
PAGE 08



BELI
PAGE 10



BUTTER
PAGE 12



CREMER
PAGE 14



KETTING
PAGE 16



KHMELINSKII
PAGE 18



KÖNIG
PAGE 20



LEGEWIE
PAGE 22



LEMKE
PAGE 24



LUKE
PAGE 26



NIEHRS
PAGE 28



RICHLY
PAGE 30



ROIGNANT
PAGE 32



ROUKOS
PAGE 34



SOSHNIKOVA
PAGE 36



ULRICH
PAGE 38



WOLF
PAGE 40



MIGUEL ANDRADE



“MANY NON-CODING RNAs EVOLVED FROM COPIES OF PROTEIN-CODING GENES”

EDUCATION

- 1994** PhD in Computational Biology, Complutense University of Madrid
- 1989** MSc in Chemistry, Complutense University of Madrid

POSITIONS HELD

- Since 2014** Adjunct Director, Institute of Molecular Biology (IMB), Mainz
Professor of Bioinformatics, Johannes Gutenberg University Mainz
- Since 2008** Affiliate Investigator, Ottawa Health Research Institute
- 2007 – 2014** Group Leader, Max Delbrück Center for Molecular Medicine, Berlin
- 2006 – 2008** Senior Scientist, Ottawa Health Research Institute
- 2003 – 2006** Scientist, Ottawa Health Research Institute
- 1998 – 2003** Staff Scientist, European Molecular Biology Laboratory (EMBL), Heidelberg
- 1996 – 1998** Postdoc, EMBL-EBI, Hinxton
- 1995 – 1996** Postdoc, National Centre for Biotechnology, Madrid
- 1994 – 1995** Postdoc, European Molecular Biology Laboratory (EMBL), Heidelberg

GROUP MEMBERS

- Gregorio Alanis-Lobato** Postdoc; since 02/2015
- Steffen Albrecht** PhD Student; since 03/2017
- Tommaso Andreani** PhD Student; since 01/2016
- Jean-Fred Fontaine** Postdoc; since 04/2014
- Jonas Ibn-Salem** PhD Student; 09/2014 – 02/2018
- Pablo Mier** Postdoc; since 01/2015
- Enrique Muro** Staff Scientist; since 04/2014
- Sweta Talyan** PhD Student; 12/2014 – 06/2018
- Katerina Taškova** Postdoc; 11/2014 – 05/2018
- Kristina Kastano** PhD Student; since 07/2018
- Michael Lenz** Postdoc; since 07/2018

OVERVIEW

Our group develops and applies methods that integrate data at different levels of molecular biology to investigate biological questions, including the function of genes and proteins and the mechanisms that control cell identity or cause disease. Our projects often overlap, both in terms of the resources and methods they use. For example, we develop data mining methods that associate keywords to therapeutic drugs, which we can then apply to the interpretation of gene expression profiles. In a different project, we created particular phylogenetic analyses of protein families that we can then use to study the evolution of the human protein interaction network. Carrying out these projects within the same group allows each project to benefit from and complement each other.

RESEARCH HIGHLIGHTS

Protein interaction networks

Geographical maps are useful because distances in them are informative. We are working to create similarly meaningful (hyperbolic) maps of protein-protein interaction networks, where the geometric properties of the map correspond to biological properties: distance to likelihood of interaction, radial distance to protein age, and angular coordinates to function. Our maps have predictive power and can be explored using an associated web server (GAPI). We evaluated the properties of disease modules (DMs) in these maps. DMs are sets of proteins associated to a single disease that agglomerate in the network because they tend to interact with common components (explaining why they associate to the same disease). Indeed, when compared to groups of proteins of similar size and connectivity but not related to the same disease, we observe that DMs form ensembles with significantly large connectivity and short spatial distances (in both the network and hyperbolic maps).

Protein sequence analysis

Following our work in the development of tools to support the study of protein evolution and function, we developed a tool (ProteinPathTracker) that aids the investigation of the evolutionary path of a protein. It is based on the retrieval of homologs (if possible, orthologues) in a series of species at increasing taxonomic distances from a central species of interest. These homologs should give information on the ancestral sequences that lead to the sequence of interest. The tool allows for testing a series of evolutionary paths, for example, from human as central species to bacteria. Beginning with a protein in the central species, the user can start a step-wise search for orthologues from species to species or, if a sequence is provided, from the closest homologue.

Figure 1. Alignment of non-overlapping fragments of a three-frame translated non-coding DNA region to a protein they are similar to. This was obtained using our similarity matrix that accounts for random DNA evolution.

Alignment of AA to DNA		
Target 1	M T N T K G K R R G T r y M F S r P F r K H G V V p L A t Y	30
Query 1	ATG ACA AAC ACA AAG GGA AAG AGG AGA GGT ACC tgg t ATG TTC TCC aag CCT TTT gga AAA CAT GGA GTT GTT cgt TTG GCC agg TAC M T N T K G K R R G T w M F S k P F g K H G V V r L A r Y	88
Target 31	M r I y K K G D i V d I K G M G T V Q K G m p H k C Y H G K	60
Query 89	ATG tga ATC tgt AAG AAA GGT GAT atg GTA ggc ATC AAG GGA ATG GGC ACT GTT CAA AAA GGT ata acc CAC agg TGT TAC CAT GGC AAA M * I c K K G D m V g I K G M G T V Q K G i t H r C Y H G K	178
Target 61	T G r V Y N V t Q h a v G I V v N k Q v k G k I L a K R I N	90
Query 179	ACT GGA agc GTT TAC AAT GTT ccc CAG tgt act gct GGC ATT GTT gca AAT gaa CAA g --- GGC atg ATT CTT gac AAG AGA ATT AAT T G s V Y N V p Q c t a G I V a N e Q --- G m I L d K R I N	263
Target 91	V r I E H I K H S k S r D S F L K r V K E N D Q K K K e a k	120
Query 264	GTG ggt ATT GAG CAC ATT AAG CAC TCT ggg AGT caa GAT AGC TTC CTG AAA cac GTG AAG GAA AAT GAT CAG AAA AAG AAG cc --- caa V g I E H I K H S g S q D S F L K h V K E N D Q K K K	349
Target 121	E K G T W v Q L K R Q p A P p R E A H f V R T n G K e P E L	150
Query 350	GAG AAG GGT ACC TGG gct CAA CTG AAG CGC CAG ett GCT CCA cgc AGA GAA GCG CAC tgt GTG AGA ACC agt GGG AAA gg CCA GAG CTG E K G T W a Q L K R Q l A P r R E A H c V R T s G K P E L	438
Target 151	L E P I P Y E F M A	160
Query 439	CTA GAA CCT ATT CCC TAT GAG TTC ATG GCA L E P I P Y E F M A	468

Transcript prediction from sequence analyses of genomes

Comparing the genomic regions of DNA that apparently do not code for proteins with their protein coding counterparts can reveal either ncRNAs that are complementary to coding RNAs, thereby potentially interacting with them, or actual coding regions that might have escaped previous detection. With this goal in mind, we developed an alignment method specifically to detect regions in non-coding DNA with similarity to protein coding genes (Figure 1). This method benefits from a substitution matrix that we use to compare three-frame translations of non-coding DNA against proteins. The similarity score is modelled for random mutations. Application of this method to human lincRNAs detected 203 transcripts with significant similarity to protein-coding genes, suggesting regulatory functions for these lincRNAs. We also contributed to a method and associated web tool (AnABlast) that detects potential coding regions in DNA by running the standard BLASTX algorithm to compare all translated frames of the query DNA sequence against a protein database. Graphical display of the accumulated hits can be used to indicate potential coding regions or their remnants.

FUTURE DIRECTIONS

We are expanding our approaches to aid the prediction of gene and protein function. For example, we are working on tools that evaluate multiple sequence alignments of proteins to find amino acids or motifs conserved in particular species. To further study low complexity protein regions, we are developing new measures for repetitiveness, which we will apply to hundreds of species whose proteomes are complete.

Finally, we are trying to explain the function of genomic regions, called Topologically Associating Domains (TADs), which are defined as having more contacts within them than with other genomic regions. To do this, we will explore the distributions of genes with different functions and breadth of expression within TADs and their dependence on the size of the TAD.

SELECTED PUBLICATIONS

Talyan S, Andrade-Navarro MA and Muro EM. (2018). Identification of transcribed protein coding remnants within lincRNAs. *Nucleic Acids Res*, 46: 8720–8729.

Alanis-Lobato G, Mier P and Andrade-Navarro MA. (2018). The latent geometry of the human protein interaction network. *Bioinformatics*, 34: 2826–2834.

Krefting J, Andrade-Navarro MA and Ibn-Salem J. (2018). Evolutionary stability of topologically associating domains is associated with conserved gene regulation. *BMC Biol*, 16: 87.

PETER BAUMANN

“TELOMERASE TIPS THE SCALES FOR AGEING AND CANCER”



EDUCATION

- 1998** PhD in Biochemistry, University College London
1994 MPhil, University of Cambridge

POSITIONS HELD

- Since 2018** Adjunct Director, Institute of Molecular Biology (IMB), Mainz
Since 2017 Alexander von Humboldt Professor, Johannes Gutenberg University Mainz
Since 2013 Professor, Kansas University Medical Center
2013 – 2018 Investigator, Howard Hughes Medical Institute, Kansas City
2013 – 2018 Priscilla Wood-Neaves Endowed Chair in the Biomedical Sciences, Stowers Institute for Medical Research, Kansas City
2013 – 2018 Investigator, Stowers Institute for Medical Research, Kansas City
2009 – 2013 Early Career Scientist, Howard Hughes Medical Institute, Kansas City
2009 – 2013 Associate Professor, Kansas University Medical Center
2009 – 2012 Associate Investigator, Stowers Institute for Medical Research, Kansas City
2004 – 2009 Assistant Professor, Kansas University Medical Center
2002 – 2008 Assistant Investigator, Stowers Institute for Medical Research, Kansas City
2000 – 2002 Research Associate, Howard Hughes Medical Institute, University of Colorado, Boulder
1998 – 2000 Wellcome Trust Travelling Research Fellow, University of Colorado, Boulder

GROUP MEMBERS

- David Ho** PhD Student; since 04/2018
Kristi Jensen Lab Manager; since 05/2018
Katja Kratz Staff Scientist; since 10/2017
Abinaya Manivannan PhD Student; since 08/2018
Aaron Odell Bioinformatician; since 01/2017
Alex Orioli PhD Student; since 01/2018
Diego Paez-Moscoso PhD Student; since 06/2014
Lili Pan Postdoc; since 02/2015
Ute Sideris Personal Assistant; since 10/2017
Chi-Kang Tseng Postdoc; since 09/2013
Hui-Fang Wang Technician; since 07/2014

OVERVIEW

Elucidating the mechanisms of chromosome end maintenance has far-reaching implications for the treatment of cancer and certain degenerative diseases characterised by the premature depletion of stem cell pools. Overall, our research goals in this area are guided by the conviction that a better understanding of telomerase biogenesis and the dynamic interactions that occur at telomeres will enable us to identify compounds that modulate telomere length. Such reagents will have therapeutic use either to limit the lifespan of tumour cells or to boost the proliferative potential of desired cell populations counteracting many of the detrimental phenotypes associated with ageing.

Our interest in chromosome dynamics and evolution has led us to a group of lizard species with fascinating attributes: They arose by interspecific hybridisation and reproduce clonally via parthenogenesis. We have revealed deviations from the normal meiotic program that permit such unisexual reproduction, and have generated cell, biological, and genomic resources to elucidate the molecular basis of clonal reproduction in a vertebrate species. Such research carries the promise of furthering our knowledge of evolution and gene regulation in the context of hybridisation, ploidy variation, and parthenogenesis.

RESEARCH HIGHLIGHTS

Telomerase biogenesis and regulation

Progressive telomere shortening eventually limits the replicative potential of cells by triggering senescence. However, replenishing telomeric sequences is a double-edged sword. It is vital for tissue homeostasis especially in long-lived species such as humans, but it also permits the continued proliferation of malignant cells. Thus, tight regulation of the enzyme telomerase is of pivotal importance. The isolation of the telomerase RNA subunit (TER1) from fission yeast in our laboratory has provided a key

tool for studying the biogenesis and regulation of the enzyme in a genetically tractable organism. This has led to a series of discoveries in telomere and RNA biology. We demonstrated that the RNA splicing machinery has a second, previously overlooked, function in RNA 3' end processing. Instead of removing an intronic sequence in a two-step process, the first transesterification reaction alone generates the mature 3' end of TER1. Our work also defined roles for several RNA chaperones in telomerase biogenesis. Most interestingly, we found that the Sm proteins and a related protein complex (LSm - Like-Sm) sequentially associate with telomerase RNA and play distinct roles in telomerase maturation. Sm and LSm proteins are members of an ancient family of RNA binding proteins that affect virtually every aspect of RNA metabolism. In contrast to the established view that they have distinct sets of RNA targets, our work revealed the first example of an RNA that requires sequential binding by the two complexes and documents specific functions for each during maturation.

While our studies on telomerase and telomeres in fission yeast have provided fundamental insights into chromosome end maintenance, they are ultimately a stepping stone towards understanding telomere maintenance in human cells. Recent work in our group uncovered roles of the cap-binding complex, the nuclear exosome targeting complex, the Trf4/Air2/Mtr4 polyadenylation complex, the exosome, and poly(A) ribonuclease in the biogenesis of human telomerase. Importantly, our studies showed that human telomerase RNA (hTR)

processing is in kinetic competition with degradation (Figure 1), an observation that hints at potential treatment options for telomerase insufficiency disorders and for delaying or reversing certain degenerative processes associated with ageing.

Mechanism of parthenogenesis

Although all-female species of whiptail lizards were already described in the early 1960s, how mature eggs are produced in the absence of fertilisation, remained an enigma until we showed that a transient doubling of chromosomes prior to the meiotic divisions produces diploid rather than haploid eggs. To gain a deeper mechanistic understanding, we have sequenced and *de novo* assembled ovarian transcriptomes and cloned key meiotic regulators. These reagents enabled us to show that the doubling of chromosomes affects only a small fraction of oocytes and have provided insights into the underlying cellular processes.




Figure 1. Telomerase biogenesis is a highly regulated multi-step process. The long precursor of human telomerase RNA is either degraded or processed into a shorter precursor that undergoes further quality control for folding and ribonucleoprotein complex assembly. Only those molecules that pass QC are processed into the mature form that is part of the active telomerase complex.

FUTURE DIRECTIONS

To gain a comprehensive understanding of human telomerase biogenesis, regulation and turnover, future studies are aimed at identifying additional factors and using biochemical and genetic means to elucidate their functions. Unravelling how telomerase is made and regulated has led us to several exciting questions: Can we modulate telomerase activity by manipulating RNA processing events? Is increasing the levels of telomerase offering a path towards the treatment of telomeropathies? Does increased telomerase activity contribute to resilience and delay processes associated with ageing?

With respect to parthenogenesis, future studies will be focused on the mechanism of chromosome doubling and the regulation of cellular functions despite the presence of multiple genomes in hybrids. These studies will inform subsequent experiments aimed at understanding the molecular events through which barriers to unisexuality are overcome. Such research has far-reaching implications for areas as diverse as evolutionary biology and agriculture.

SELECTED PUBLICATIONS

Páez-Moscoso DJ, Pan L, Sigauke RF, Schroeder MR, Tang W and Baumann P. (2018). Pof8 is a La-related protein and a constitutive component of telomerase in fission yeast. *Nat Commun*, 9: 587.

Newton AA, Schnittker RR, Yu Z, Munday SS, Baumann DP, Neaves WB and Baumann P. (2016). Widespread failure to complete meiosis does not impair fecundity in parthenogenetic whiptail lizards. *Development*, 143: 4486–4494.

Tseng CK, Wang HF, Burns AM, Schroeder MR, Gaspari M and Baumann P. (2015). Human telomerase RNA processing and quality control. *Cell Rep*, 13: 2232–2243.

PETRA BELI

“WE STUDY UBIQUITIN AND PHOSPHORYLATION-DEPENDENT PROCESSES IN THE DNA DAMAGE RESPONSE”



EDUCATION

- 2011** PhD in Biology, Goethe University Frankfurt
2007 MSc in Molecular Biology, University of Zagreb

POSITIONS HELD

- Since 2013** Emmy Noether Group Leader, Institute of Molecular Biology (IMB), Mainz
2010 – 2013 Postdoc, Novo Nordisk Foundation Center for Protein Research, University of Copenhagen

GROUP MEMBERS

- Irem Baymaz** Postdoc; since 11/2016
Marina Borisova PhD Student; 12/2013 – 07/2018
Francesca Conte Research Assistant; since 09/2018
Sofia Lobato Gil Postdoc; since 10/2018
Jan Heidelberg PhD Student; since 04/2014
Thomas Juretschke PhD Student; since 09/2016
Thorsten Mosler PhD Student; since 04/2017
Matthias Ostermaier PhD Student; since 05/2015
Andrea Voigt Lab Manager; since 01/2014
Juanjuan Wang PhD Student; since 08/2016

OVERVIEW

The genome integrity of living organisms is challenged by by-products of cellular metabolism and external factors such as ultraviolet (UV) light and ionising radiation. To maintain genome integrity, mammalian cells have evolved elaborate mechanisms, jointly known as the DNA damage response (DDR), that regulates DNA repair and cell cycle checkpoints. Germline or somatic mutations in DDR genes that affect the cellular capacity to respond to DNA damage result in the accumulation of mutations. This consequently leads to syndromes that manifest as premature ageing and the development of cancer. Protein phosphorylation mediated by Ataxia-telangiectasia mutated kinase (ATM), Ataxia telangiectasia and Rad3 related (ATR) and DNA-dependent protein kinases (DNA-PKcs) play essential roles during the cellular response to DNA damage. We are employing quantitative mass spectrometry (MS) based proteomics to identify and characterise phosphorylation-and ubiquitin-dependent signalling networks that regulate the DDR in human cells.

RESEARCH HIGHLIGHTS

Recent studies revealed that exposure of human cells to UV light affects transcription, splicing and translation. These findings raised a number of questions: Are these changes a regulated process that is important for genome stability maintenance? If so, which signalling pathways and molecular mechanisms link DNA damage with the regulation of RNA metabolism? In addition to ATR-Chk1, studies have shown that p38 and JNK MAPK signalling is activated after UV light, however, the substrates and functions of these pathways remained poorly understood.

We combined kinase inhibition and quantitative phosphoproteomics to analyse kinase-dependent signalling in response to UV light. We defined the cellular phosphorylation events dependent on canonical DNA damage signalling mediated by the ATR-Chk1

and the p38 MAPK pathways and determined the functional contributions of these pathways to the UV light-induced DDR. Whereas ATR primarily phosphorylates proteins that function in DNA repair and cell cycle regulation, the p38-MK2 signalling axis phosphorylates a multitude of RNA-binding proteins (RBPs). We identified 138 phosphorylation sites on 122 proteins to be dependent on p38-MK2 signalling. We showed that p38-MK2-dependent phosphorylation of cellular proteins triggers the recruitment of 14-3-3 dimers, thus providing a general mechanism that rapidly regulates RBPs after UV light exposure.

Furthermore, we demonstrated that the negative elongation factor (NELF) complex, which plays an essential role in promoting RNA polymerase II (RNA pol II) promoter-proximal pausing, is a substrate of p38-MK2. Biochemical studies and X-ray crystallography revealed that NELF complex phosphorylation leads to the recruitment of 14-3-3 to the RNA binding subunit NELFE. We

employed SILAC-based MS to systematically probe the composition of chromatin after UV light exposure as well as to identify p38-dependent changes in protein recruitment to or dissociation from chromatin. Phosphorylation and subsequent interaction of the NELF complex with 14-3-3 promotes its rapid release from chromatin, which correlates with RNA pol II elongation.

Taken together, the results of our study provided insights into the function of the p38-MK2 signalling pathway in the regulation of RBPs and revealed a phosphorylation-dependent mechanism that promotes RNA pol II elongation and DNA repair in response to UV light (Figure 1). We anticipate that the provided datasets of UV light-induced phosphorylation sites and p38-dependent 14-3-3 interactions will enable further studies focusing on the functions of the p38-MK2 pathway in the regulation of different RNA metabolic processes after UV light exposure.




Figure 1. Model for the UV light-induced regulation of transcriptional elongation through the p38-MK2-NELF signalling axis. p38 and its downstream effector kinase MK2 phosphorylate a multitude of RBPs after UV light. Phosphorylation of NELFE results in 14-3-3 binding and release of the NELF complex from chromatin that is accompanied by RNA pol II elongation.

FUTURE DIRECTIONS

Posttranslational modifications of proteins by phosphorylation and ubiquitylation play important regulatory roles in the cellular response to DNA damage. However, the complete picture of posttranslational regulatory events that control the DNA damage response and its interplay with other cellular processes is missing. We will employ quantitative proteomics, biochemistry

and cell biology to identify and characterise phosphorylation and ubiquitin-dependent processes that govern the DNA damage response. Furthermore, we aim to develop proteomics methods that will enable us to obtain an unbiased view of the chromatin proteome at sites of DNA damage.

SELECTED PUBLICATIONS

Borisova ME, Voigt A, Tollenaere MAX, Sahu SK, Juretschke T, Kreim N, Mailand N, Choudhary C, Bekker-Jensen S, Akutsu M, Wagner SA and Beli P. (2018). p38-MK2 signaling axis regulates RNA metabolism after UV-light-induced DNA damage. *Nat Commun*, 9: 1017.

Heidelberg JB, Voigt A, Borisova ME, Petrosino G, Ruf S, Wagner SA and Beli P. (2018). Proteomic profiling of VCP substrates links VCP to K6-linked ubiquitylation and c-Myc function. *EMBO Rep*, 19: e44754.

Wagner SA, Oehler H, Voigt A, Dalic D, Freiwald A, Serve H and Beli P. (2016). ATR inhibition rewires cellular signalling networks induced by replication stress. *Proteomics*, 16: 402–416.

FALK BUTTER

“WE APPLY QUANTITATIVE PROTEOMICS APPROACHES TO STUDY EVOLUTION AND GENOME STABILITY”



EDUCATION

- 2010** PhD in Biochemistry, Ludwig Maximilian University (LMU), Munich
- 2006** Diploma in Biochemistry, University of Leipzig

POSITIONS HELD

- Since 2013** Group Leader, Institute of Molecular Biology (IMB), Mainz
- 2010 – 2013** Postdoc, Max Planck Institute for Biochemistry, Martinsried

GROUP MEMBERS

- Alina Bluhm** PhD Student; since 01/2014
- Hanna Braun** PhD Student; since 08/2015
- Núria Casas Vila** PhD Student; 11/2013 – 12/2018
- Sabrina Dietz** PhD Student; since 01/2015
- Albert Fradera Sola** PhD Student; since 03/2018
- Teresa Laguna** Postdoc; 03/2016 – 04/2018
- Michal Levin** Postdoc; since 09/2018
- Katarina Luko** PhD Student; since 12/2017
- Tony May** Master's Student; 11/2017 – 09/2018
- Emily Nischwitz** PhD Student; since 09/2018
- Merve Öztürk** PhD Student; since 11/2015
- Lara Perez** PhD Student; since 10/2015
- Marion Scheibe** Postdoc; since 06/2013
- Marian Scherer** Bachelor's Student; since 11/2017
- Vivien Schoonenberg** PhD Student; since 02/2018
- Nikenza Viceconte** Postdoc; since 01/2018

OVERVIEW

Mass spectrometry is a powerful tool for studying proteins in an unbiased and global manner. The current improvements in identification accuracy, sample throughput, and data analysis allow the streamlined application of proteomics in answering diverse biological questions. Our group applies quantitative approaches, such as label-free quantitation (LFQ), reductive demethylation (DML) or stable isotope labelling with amino acids in cell culture (SILAC), which enables us to directly compare thousands of proteins in complex mixtures. These technologies allow us to study changes in protein expression and are also applied in interactomics to identify specific interactions of proteins within a vast number of background binders. We apply mass spectrometry in several biological areas to advance our knowledge of cellular processes.

RESEARCH HIGHLIGHTS

Phylointeractomics reveals evolutionary changes in protein binding

We developed a new experimental workflow for comparative evolutionary biology, termed “phylointeractomics”. Built upon the ability of mass spectrometry-based proteomics to identify proteins from sequence information, it is a perfect technique for cross-species comparison. In phylointeractomics, we interrogate a bait of interest with the proteome of evolutionarily related species in a systematic manner to uncover similarities and differences in protein binding. In a first application, we studied the telosome of 16 different vertebrate species ranging from zebrafish to human, which span a timeframe of 450 million years of evolution. While the telomeric sequence in vertebrates is a conserved TTAGGG repeat, there are some known variations of the interacting proteins, e.g. a Pot1 gene duplication in the rodent lineage and absence of TIN2 in several bird genomes. In our phylointeractomics screen, we recapitulated these evo-

lutionary differences for the shelterin complex and additionally uncovered that, in contrast to predictions, not all homologues of TRF1, a direct TTAGGG-repeat binding subunit of the complex, associated with our telomeric baits. Using recombinant expressed TRF1 DNA-binding domains of even more vertebrate species, we could locate a gain-of-binding event at the branch point of the therian lineage, where mammals and marsupials diverged from monotremes such as the platypus. While TRF1 is present in most vertebrates, it seems to have obtained its telomeric function only later during vertebrate diversification. By exchange of selected amino acid residues in the platypus TRF1-DNA binding domain, we could recreate a gain-of-binding switch *in vitro* recapitulating a possible evolutionary scenario. Our phylointeractomics study therefore underscores that sequence homologues, as determined by phylogenomics, do not necessarily need to equate to functional homology.




Figure 1. Microscopy-based approaches to study telomere-binding proteins. We investigated the candidate ZBTB10 by comparing U2OS WT and knock-out clones that were labelled with BrdU:BrdC and arrested in metaphase. Telomeres were stained with two FISH probes to quantify telomere sister chromatid exchange (left panel). Immunofluorescence staining of PML protein (ALT-associated PML body marker) and TRF2 (telomere marker) to determine co-localisation with ZBTB10 (upper right panel). Imaging of telomere-induced DNA damage foci with a TTAGGG FISH probe and 53BP1 antibody (lower right panel).

Characterisation of new telomeric proteins

We use quantitative interactomics to identify new telomeric proteins. Apart from HOT1, we recently reported that the zinc finger protein ZBTB48 is a telomeric protein in mammals. The extension of our workflow to other model species resulted in the identification of novel telomere binding proteins in *Trypanosoma brucei*. We characterised one of these candidates, which we termed TelAP1 as a dynamic telomere binding protein that is differentially expressed between the two life cycle stages of the parasite.

Systems approaches to study developmental gene regulation

To study proteome dynamics during development, we generated two large developmental proteomic datasets of *Drosophila melanogaster*: a full life cycle dataset encompassing 15 different time points and a highly temporally-resolved proteome of its embryogenesis. As both datasets match the previously published modENCODE developmental transcriptome, we systematically compared developmental transcriptome and proteome expression, showing that for selected cases protein stability is the major determinant of protein levels. Additionally, we identified maternally loaded proteins, uncovered peptides originating from small open reading frames in lncRNAs and resurrected the pseudogene (Cyp9f3). The data is available to the research community via our web interface (www.butterlab.org/flydev). Further, we teamed up with the Legewie group to use our large-scale transcriptomics and proteomics datasets to investigate posttranslational gene regulation by mathematical modelling.

FUTURE DIRECTIONS

We will continue to apply quantitative proteomics to diverse biological questions with a focus on differentiation, epigenetics, development and evolution. To this end, we are currently improving several parts of the proteomics and interactomics workflow

established during the last few years in our group. Combining omics studies with classical biology, we are at the moment characterising novel telomeric proteins in diverse model species and investigating gene regulation in several eukaryotes.

SELECTED PUBLICATIONS

Casas-Vila N*, Bluhm A*, Sayols S*, Dinges N, Dejung M, Altenhein T, Kappei D, Altenhein B, Roignant JY and Butter F. (2017). The developmental proteome of *Drosophila melanogaster*. *Genome Res*, 27: 1273–1285.

* indicates joint contribution # indicates joint correspondence

Kappei D*, Scheibe M*, Paszkowski-Rogacz M, Bluhm A, Gossmann TI, Dietz S, Dejung M, Herlyn H, Buchholz F*, Mann M* and Butter F*. (2017). Phylointeractomics reconstructs functional evolution of protein binding. *Nat Commun*, 8: 14334.

Reis H, Schweps M, Dietz S, Janzen CJ and Butter F. (2018). TelAP1 links telomere complexes with developmental expression site silencing in African trypanosomes. *Nucleic Acids Res*, 46: 2820–2833.

CHRISTOPH CREMER

“EPIGENETICS IS
CONTROLLED BY
SPATIAL CONSTRAINTS
AT THE NANOSCALE”



EDUCATION

- 1983** Habilitation in General Human Genetics and Experimental Cytogenetics, University of Freiburg
- 1976** PhD in Biophysics and Genetics, University of Freiburg
- 1970** Diploma in Physics, Ludwig Maximilian University (LMU), Munich

POSITIONS HELD

- Since 2015** Research Associate, Max Planck Institute for Chemistry, Mainz
- Since 2013** Honorary Professor, Johannes Gutenberg University Mainz
- Since 2011** Group Leader, Institute of Molecular Biology (IMB), Mainz
- Since 2005** Director, Cooperation Unit Biophysics, Institute for Pharmacy and Molecular Biotechnology, University of Heidelberg
- 2005 – 2007** Deputy Director, Kirchhoff Institute of Physics, University of Heidelberg
- 1983 – 2011** Professor of Applied Optics & Information Processing, University of Heidelberg
- 1983 – 1999** Managing/Deputy Director, Institute of Applied Physics I, University of Heidelberg
- 1970 – 1983** Staff Scientist, Institute of Human Genetics, University of Freiburg

GROUP MEMBERS

- Shih-Ya Chen** PhD Student; since 05/2016
- Marton Gelleri** Postdoc; since 05/2018
- Maria Contreras Gerenas** Postdoc; since 02/2018
- Xiaomin Liu** Postdoc; since 11/2018
- Renata Pandolfo** PhD Student; since 11/2017
- Florian Schock** PhD Student; since 03/2014
- Felix Schreiber** Bachelor's Student; since 07/2018
- Fuguang Zhao** PhD Student; since 11/2018

OVERVIEW

A complete mechanistic understanding of gene regulation in space and time requires a detailed insight into the spatial constraints of gene activity control in the cell nucleus, along with sequence and biochemical information. For this, it is necessary to enhance the resolution of the spatial DNA distribution and other relevant molecules substantially beyond the conventional resolution level. This has become possible by various super-resolution light microscopy (SRM) approaches. SRM of the nuclear genome nanostructure is expected to contribute to fundamental questions including how does the complex organisation of chromosome territories in the mammalian cell nucleus arise, and how is this organisation related to epigenetic gene regulation and its dynamics? We have shown that Single Molecule Localisation Microscopy (SMLM), a type of SRM, is particularly useful for enhanced analysis of the cell nucleus. In 2018, we applied a recently developed SMLM approach to nuclear genome nanostructure imaging, designated DNA Structure Fluctuation Assisted Binding Activated Localisation Microscopy (fBALM), to analyse the mammalian cell nucleus in three dimensions at unprecedented resolution (optical and structural 3D single molecule resolution of few tens of nm). Present application fields include neurobiology (genome nanostructure of neurons) and human cancer tissues.

RESEARCH HIGHLIGHTS

In the last year, we focused on the following topics: What are the differences in DNA content between the low density (active) and the high density (inactive) compartments? What is the spatial nanoscale distribution of such small chromatin domains? How many nucleosomes do they contain? What is the variability of such nucleosome clusters? Information on these questions is essential for a quantitative, mechanistic and dynamic understand-

ding of epigenetic gene regulation. As such, we have applied a new technique (fBALM), developed in our group, for localisation microscopy of nuclear DNA at single molecule resolution. This substantially enhanced the structural resolution, down to a few tens of nm in 3D. The fBALM localisation microscopy confirmed that the nuclear genome is organised into distinct, presumably inactive, high-density domains with a size substantially below 100 nm in diameter, and, presumably active, low-density compartments (Figure 1). It was possible to obtain estimates of the absolute number of nucleosomes (DNA content) in such nanoclusters. New software packages were developed for an improved quantitative nanocluster analysis. To further enhance resolution, we completed the construction of a new 3D Single Molecule Localisation Microscope (3D-SMLM) in the last year. Using a spatial light modulator system, it is now possible to illuminate specific cells or intracellular regions and to enhance the homogeneity of the illumination. In addition, the 3D localisation mode of the new 3D-SMLM system allowed measuring up to 5.5 million different DNA sites throughout an entire three-dimensional nucleus, with a precision of about 10 nm. We also combined the fBALM method with Single Molecule Localisation Analysis using immunostaining. In addition, we

constructed and implemented a multiplex micropump system; this system now makes it possible to prolong fBALM experiments up to around 10 hours. This will allow us to register up to 1 million image frames and thus map the nuclear 3D DNA distribution at unprecedented resolution (up to 30 million DNA sites/nucleus corresponding to 1 DNA position per nucleosome).

In collaboration with wet lab scientists, the new fBALM approaches were used to address a variety of research questions such as studying genome nanostructure in tissue sections from human colon cancer and in mouse neuronal cells. Our first experiments with Light Sheet Microscopy allowed us to identify almost all nuclei in thick mouse brain tissue sections. Finally, in a collaboration with the Max Planck Institute for Chemistry, allergy-related membrane clusters (TLR4) and the translocation of NfκB to the nucleus were studied.




Figure 1. The image on the left shows the fBALM image of a single HeLa cell nucleus. The image consists of approximately 4 million single molecule localisations each of them corresponding to individual DNA sites. The inspection of small insets in such images (right) reveals the chromatin nanostructure.

FUTURE DIRECTIONS

In addition to the continuation of the projects described above, the SRM instrumentations and the imaging and data evaluation methods developed will be applied in further collaborative projects. In one such example, in collaboration with the MPI for Polymer Research, an extremely fast switchable laser system will be adapted for SRM to facilitate multiplexed localisation microscopy.

The long-term perspectives of SRM of the nuclear landscape will be the integration of the collected collaborative data into

a quantitative, mechanistic, predictive and dynamic space-time model of functional nuclear organisation and its integration into molecular biology approaches. Using super-resolution techniques, it should become possible to directly measure transcription induced condensation changes in space and time on the level of very small, individual nuclear domains in single nuclei and to study the nanostructural consequences of chromatin modifiers, e.g. histone/DNA methylases and demethylases.

SELECTED PUBLICATIONS

Szczurek A, Klewes L, Xing J, Gourram A, Birk U, Knecht H, Dobrucki JW, Mai S and Cremer C. (2017). Imaging chromatin nanostructure with binding-activated localisation microscopy based on DNA structure fluctuations. *Nucleic Acids Res*, 45: e56

Kirmes I, Szczurek A, Prakash K, Charapitsa I, Heiser C, Mushev M, Schock F, Fornalczyk K, Ma D, Birk U, Cremer C and Reid G. (2015). A transient ischemic environment induces reversible compaction of chromatin. *Genome Biol*, 16: 246.

Prakash K, Fournier D, Redl S, Best G, Borsos M, Ketting R, Tachibana-Konwalski K, Cremer C and Birk U. (2015). Super-resolution imaging reveals structurally distinct periodic patterns of chromatin along pachytene chromosomes. *Proc Natl Acad Sci*, 112: 14635–14640.

RENÉ KETTING

“GERM CELLS CRITICALLY
DEPEND ON SMALL RNA
ACTIVITY”



EDUCATION

- 2000** PhD in Molecular Biology, Netherlands Cancer Institute, Amsterdam
- 1994** MSc in Chemistry, University of Leiden

POSITIONS HELD

- Since 2012** Scientific Director, Institute of Molecular Biology (IMB), Mainz
Professor, Faculty of Biology, Johannes Gutenberg University Mainz
- 2015 – 2017** Executive Director, Institute of Molecular Biology (IMB), Mainz
- 2010 – 2013** Professor of Epigenetics in Development, University of Utrecht
- 2005 – 2012** Group Leader, Hubrecht Institute, Utrecht
- 2000 – 2004** Postdoc, Hubrecht Institute, Utrecht
- 2000** Postdoc, Cold Spring Harbor Laboratories

GROUP MEMBERS

- Miguel Almeida** PhD Student; since 06/2013
- Walter Bronkhorst** Postdoc; since 01/2015
- Alessandro Consorte** PhD Student; since 01/2018
- Antonio Domingues** Postdoc; since 12/2014
- Yasmin El Sherif** Lab Manager; since 10/2014
- Svena Hellmann** Technician; since 11/2016
- Monika Kornowska** Animal Caretaker; since 03/2015
- Saskia Krehbiel** Animal Caretaker; since 05/2012
- Regina Otto** Personal Assistant; since 09/2015
- Maria Placentino** PhD Student; since 11/2013
- Nadezda Podvalnya** PhD Student; since 01/2018
- Stefan Redl** PhD Student; since 04/2013
- Ricardo Rodrigues** PhD Student; since 12/2012
- Elke Roovers** PhD Student; since 07/2013
- Jan Schreier** PhD Student; since 05/2015
- Nadine Wittkopp** Postdoc; since 11/2012

OVERVIEW

The major focus of my lab is on gene regulation by small RNA molecules acting in RNAi-related pathways. Since their discovery at the start of the 21st century, many different RNAi-related pathways have been identified. It is now evident that although all of these pathways depend on proteins from the Argonaute family, each pathway has its own unique characteristics and effects on gene expression. These can range from relatively minor effects on translation (in the case of miRNAs) to full-blown shutdown of loci at the transcriptional level (piRNAs). We focus on mechanisms related to piRNA and siRNA biology, two species of small RNAs that are particularly abundant in, and important for, the germline. These small RNA pathways have a major role in maintaining genome integrity through controlling the activity of transposable elements. We use zebrafish and *C. elegans* as model systems to understand the molecular mechanisms governing these pathways and to understand how these pathways contribute to normal development. Questions including, how do small RNA pathways distinguish transposable elements from regular genes, how are these pathways organised at a sub-cellular level, and how can small RNA populations be inherited across generations are at the heart of our research.

RESEARCH HIGHLIGHTS

Flexible usage of a conserved protein in diverse small RNA pathways

In studying the factors required for small RNA pathways in different organisms, we found that the proteins acting in these pathways tend to evolve quickly. In many cases, factors that appear to be very species-specific are essential for such pathways. This likely reflects the biological niche in which these pathways operate: the control of endogenous parasites. We studied one of the few strongly conserved proteins that has been described

to act in the Piwi pathway of the fruit fly, GTSF1. Strikingly, we found that even though this protein is conserved at the sequence level, its molecular function is not. This protein has been found to trigger transcriptional silencing in flies, whereas we find it stimulates small RNA biogenesis in *C. elegans*. This protein allows the assembly of a larger molecular machinery that drives small RNA biogenesis. This principle may well apply to the fly, where GTSF1 might drive the assembly of a larger complex that instigates chromatin modification. This research emphasises that we can learn a lot from studying one factor in different organisms. Whereas one would have intrinsically coupled GTSF1 to chromatin biology based on the data from the fruitfly, our work, which was published this year in *The EMBO Journal*, shows that instead, it acts as a protein that allows the assembly of larger protein, or protein-RNA complexes.

Control of phase separation during germ cell development

We have been studying the effect of a piRNA-pathway component, named Tdrd6, on germ cell formation in zebrafish. Maternal loss of this protein leads to less efficient germ cell specification, but the molecular reasons behind this phenotype have remained unclear. We have now been able to show, using single-cell RNA sequencing, that Tdrd6 plays a role in ensuring every germ cell specified receives a certain ratio of mRNA species from different germ cell-specifying genes. Tdrd6 is itself present in granules that also bind such mRNA species, as can be visualised by single molecule FISH studies on zebrafish embryos. Tdrd6 drives the loading

of these transcripts into germ cells by helping the fusion of small RNA-protein aggregates into bigger units. This enables larger aggregates of these germ cell specifying mRNAs, also known as germ plasm, to be loaded into the future germ cells. Finally, we found that Tdrd6 performs this function by controlling the phase separation behaviour of a core germ plasm component named Bucky Ball (Figure 1). In absence of Tdrd6, Bucky Ball aggregates into amyloid-like structures that are no longer functional. This work demonstrates how, within a living animal, phase separation can be controlled and what the implications are if this control fails. This work was published this year in *Developmental Cell*.




Figure 1. A Bucky Ball aggregate surrounded by Tdrd6a. Tdrd6a binds Bucky Ball and affects its aggregation behaviour. Tdrd6a is a very dynamic protein, while Bucky Ball is rather static. Tdrd6a increases Bucky Ball mobility. The image shows an aggregate in a 4 cell stage zebrafish embryo. These aggregates are required to specify germ cells.

FUTURE DIRECTIONS

Our future work will continue to mechanistically unravel the molecular pathways that are steered by small RNA guides. We are performing a genetic screen in order to identify novel factors and are increasingly using biochemical approaches to begin to describe their mechanisms on a more molecular level. Both *C. elegans* and zebrafish will continue to play important roles in these studies. For instance, we will focus on a protein complex newly identified in our laboratory that is essential for

the generation of piRNAs. This complex is also required for an as yet unidentified but essential pathway that acts during early embryogenesis. We will also further study phase separation and how it can be controlled *in vivo*. We recently identified a novel phase-separated structure that is specific to sperm and is essential for paternal inheritance of small RNA mediated gene control. We aim to unravel the molecular details behind this new intriguing structure and how it is controlled.

SELECTED PUBLICATIONS

Kaaij LJ, Mokry M, Zhou M, Musheev M, Geeven G, Melquiond ASJ, de Jesus Domingues AM, de Laat W, Niehrs C, Smith AD and Ketting RF. (2016). Enhancers reside in a unique epigenetic environment during early zebrafish development. *Genome Biol*, 17: 146.

de Albuquerque BFM, Placentino M and Ketting RF. (2015). Maternal piRNAs are essential for germline development following *de novo* establishment of endo-siRNAs in *Caenorhabditis elegans*. *Dev Cell*, 34: 448–456.

de Albuquerque BFM, Luteijn MJ, Cordeiro Rodrigues RJ, van Bergeijk P, Waaijers S, Kaaij LJ, Klein H, Boxem M and Ketting RF. (2014). PID-1 is a novel factor that operates during 21U-RNA biogenesis in *Caenorhabditis elegans*. *Genes Dev*, 28: 683–688.

ANTON KHMELINSKII

“WE AIM TO DECIPHER
CELLULAR MECHANISMS
OF PROTEIN QUALITY
CONTROL”



EDUCATION

- 2010** PhD in Biology, University of Heidelberg
- 2005** Licenciatura degree in Biochemistry, University of Lisbon

POSITIONS HELD

- Since 2018** Group Leader, Institute of Molecular Biology (IMB), Mainz
- 2013** Visiting Scientist, Donnelly Centre for Cellular and Biomolecular Research, University of Toronto
- 2011 – 2017** Postdoc, Center for Molecular Biology (ZMBH), University of Heidelberg
- 2011 – 2016** Visiting Scientist, European Molecular Biology Laboratory (EMBL), Heidelberg
- 2010 – 2011** Postdoc, European Molecular Biology Laboratory (EMBL), Heidelberg

GROUP MEMBERS

- Iryna Charapitsa** Postdoc; 04/2018 – 12/2018
- Jia Jun Fung** PhD Student; since 01/2018
- Elena Ivanova** Research Assistant; since 07/2018
- Zhaoyan Li** PhD Student; since 12/2018
- Rocio Nieto-Arellano** Postdoc; since 04/2018
- Anke Salzer** Lab Manager; since 01/2018
- Simone Sneed** Technician; since 10/2018

OVERVIEW

The functional state of a cell is ultimately defined by the state of its proteome, i.e. abundance, localisation, turnover and mobility of all proteins and their organisation in complexes and organelles. Numerous cellular systems contribute to proteome homeostasis through prevention, detection and removal of abnormal proteins. Selective protein degradation by the ubiquitin-proteasome systems plays a key role in proteome turnover and quality control. When degradation is not possible, the impact of abnormal proteins can be minimised through their asymmetric partitioning during cell division. Despite the activity of such systems, proteome homeostasis declines with ageing and in numerous diseases, resulting in accumulation of abnormal proteins and loss of cell functionality.

Working in yeast and human cells, we aim to systematically identify substrates for the various components of the ubiquitin-proteasome system and explore the functions of this system in replicative ageing and genome stability. We are using genetics and proteomic approaches that exploit fluorescent timers to follow protein trafficking, inheritance and degradation down to subcellular resolution. Our goals are to understand how protein quality control is coordinated with protein synthesis, folding and trafficking, to elucidate how cells recognise abnormal proteins and how they adapt to challenges in proteome homeostasis.

RESEARCH HIGHLIGHTS

Selective protein degradation is involved in most cellular processes and contributes to proteome homeostasis through the removal of unnecessary or abnormal proteins. The ubiquitin-proteasome system (UPS) plays a key role in selective protein degradation, whereby a cascade of ubiquitin-activating (E1), ubiquitin-conjugating (E2) and ubiquitin-protein ligase (E3) enzymes mark proteins with polyubiquitin chains for degradation by the protea-

some. Deubiquitinating enzymes, which remove ubiquitin marks and replenish the pool of free ubiquitin, are involved at various stages of the targeting and degradation processes. Despite the central role of the UPS in protein degradation and its association with various diseases and ageing, many UPS components remain poorly characterised and our understanding of specificity in the UPS is inadequate.

We made some progress in identifying substrates and functions for various UPS components in the budding yeast *Saccharomyces cerevisiae*. Using a proteomic approach that relies on fluorescent timers as reporters of protein turnover, we were able to examine the impact of inactivating individual UPS components on the yeast proteome. This led us to the discovery of a protein quality control pathway that operates specifically at the inner nuclear membrane (INM). This pathway involves the Asi E3 ligase and appears to have a dual function: to control the abundance of specific INM-resident proteins, in a manner resembling classical endoplasmic reticulum-associated protein degradation pathways, and to maintain the identity of the INM, through recognition and targeting for degradation proteins that mislocalise to the INM.

In addition, we developed methods to facilitate further studies of the yeast proteome and address questions of specificity in the ubiquitin-proteasome system. Genome-wide libraries of strains in which each open reading frame (ORF) is fused to the same tag at its endogenous chromosomal locus are very useful tools to study the yeast proteome. But their construction is laborious and expensive, hindering the use of new tags in proteomic studies. To address these limitations, we developed the SWAp-Tag (SWAT) approach for high-throughput tagging of yeast ORFs (Figure 1). Using SWAT we are now able to endogenously tag 86% of yeast ORFs with virtually any tag in approximately three weeks, greatly expanding the possibilities of proteome-wide studies in yeast. Furthermore,

we contributed to the development of multiplexed protein stability (MPS) profiling, a method to study protein degradation signals. These signals, also known as degrons, are typically short linear motifs. Proteins with exposed degrons are recognised by cellular quality control machinery and sent for degradation. With MPS profiling, it is possible to identify and dissect degrons by examining how the turnover of a reporter protein is influenced by fusing it to thousands of different protein fragments in parallel. This method will help us understand specificity in the ubiquitin-proteasome system.




Figure 1. High-throughput protein tagging with SWAT. Individual ORFs are marked with an acceptor module in SWAT strains. New strains can be derived from SWAT strains by swapping the acceptor module for practically any tag provided on a donor plasmid. Swapping is induced by introducing double-strand breaks in the acceptor module and the donor plasmid, and involves sequentially growing yeast strains on different selective media. Double-strand break repair by homologous recombination leads to the replacement of the acceptor module by the tag.

FUTURE DIRECTIONS

We will continue our systematic search for substrates of E3 ubiquitin ligases in yeast by combining our reporter-based screens with mass spectrometry and expand this search to human cells. We will also apply genetic and proteomic approaches to identify redundancies in the ubiquitin-proteasome system and in this way,

find substrates for overlapping degradation pathways. Finally, we will build on our work on INM-associated protein degradation to understand how cells recognise mislocalised proteins. We are eager to test the importance of such quality control pathways under stress and during the ageing process.

SELECTED PUBLICATIONS

Kats I, Khmelinskii A, Kschonsak M, Huber F, Knieß RA, Bartosik A and Knop M. (2018). Mapping degradation signals and pathways in a eukaryotic N-terminome. *Mol Cell*, 70: 488-501.

* indicates joint contribution
indicates joint correspondence

Meurer M, Duan Y, Sass E, Kats I, Herbst K, Buchmüller BC, Dederer V, Huber F, Kirrmaier D, Štefl M, Van Laer K, Dick TP, Lemberg MK, Khmelinskii A*, Levy ED* and Knop M*. (2018). Genome-wide C-SWAT library for high-throughput yeast genome tagging. *Nature Methods*, 15: 598-600.

Khmelinskii A*, Blaszczyk E*, Pantazopoulou M, Fischer B, Omnis DJ, Le Dez G, Bossard A, Gunnarsson A, Barry JD, Meurer M, Kirrmaier D, Boone C, Huber W, Rabut G, Ljungdahl PO and Knop M. (2014). Protein quality control at the inner nuclear membrane. *Nature*, 516: 410-413.

JULIAN KÖNIG

“WE AIM TO CRACK
THE SPLICING CODE”



EDUCATION

- 2008** PhD in Biology, Max Planck Institute for Terrestrial Microbiology & Philipps University, Marburg
- 2003** Diploma in Biology, Ludwig Maximilian University (LMU), Munich

POSITIONS HELD

- Since 2013** Group Leader, Institute of Molecular Biology (IMB), Mainz
- 2008 – 2013** Postdoc, MRC Laboratory of Molecular Biology, Cambridge

GROUP MEMBERS

- Andreas Buchbender** PhD Student; since 04/2017
- Simon Braun** PhD Student; 11/2013 – 09/2018
- Stefanie Ebersberger** Postdoc; since 04/2014
- Heike Hänel** Lab Manager; since 11/2013
- Andrea Hildebrandt** PhD Student; since 04/2014
- Nadine Körte** PhD Student; since 11/2018
- Holger Mutter** Student Assistant; since 07/2018
- Laura Schulz** PhD Student; since 04/2017
- Reymond Sutandy** PhD Student; 12/2013 – 11/2018
- Kerstin Tretow** PhD Student; since 12/2018

OVERVIEW

Posttranscriptional regulation of gene expression at the level of splicing and translation plays a critical role in development and tissue identity. Since these processes are often implicated in disease, their detailed investigation is fundamental to our understanding of human biology in general and disease processes in particular.


The information in the RNA sequence and how it is read and interpreted by RNA-binding proteins (RBPs) is commonly referred to as the “splicing code”. Cracking this code remains a long-standing goal of RNA biologists. To further advance these efforts, our group uses functional genomics approaches to dissect the underlying regulatory networks. We developed new technologies to study the role of cis-regulatory RNA sequence elements and protein-RNA interactions at an unprecedented level of detail and resolution. Currently, we are addressing the following key questions: What is the role of cis-regulatory elements in alternative splicing? How do proteins assemble into functional ribonucleoprotein (RNP) complexes? How do RNPs act in gene expression quality control?

RESEARCH HIGHLIGHTS

Decoding a cancer-relevant splicing decision in the RON proto-oncogene using high-throughput mutagenesis

Mutations causing aberrant splicing are frequently implicated in human diseases including cancer. Together with Kathi Zarnack’s group at Goethe University Frankfurt and Stefan Legewie’s group at IMB, we established a high-throughput screen of randomly mutated minigenes (Figure 1) to decode the cis-regulatory landscape that determines alternative splicing of exon 11 in the proto-oncogene MST1R (RON). Mathematical modelling of splicing kinetics enabled us to identify more than 1000

mutations affecting RON exon 11 skipping, which corresponds to the pathological isoform RON Δ 165. Importantly, the measured effects correlated with RON alternative splicing in cancer patients bearing the same mutations. Moreover, we highlighted heterogeneous nuclear ribonucleoprotein H (HNRNPH) as a key regulator of RON splicing in healthy tissues and cancer. Using iCLIP and synergy analysis, we pinpointed the functionally most relevant HNRNPH binding sites and demonstrated how cooperative HNRNPH binding facilitates a splicing switch of RON exon 11. Our results thereby offer insights into splicing regulation and the impact of mutations on alternative splicing in cancer.



In vitro iCLIP-based modelling uncovers how the splicing factor U2AF2 relies on regulation by co-factors

Alternative splicing generates distinct mRNA isoforms and is crucial for proteome diversity in eukaryotes. The RBP U2AF2 is central to splicing decisions, as it recognises 3' splice sites and recruits the spliceosome. Together with Stefan Legewie's group at IMB, we established *in vitro* iCLIP experiments, in which recombinant RBPs are incubated with long transcripts, to study how U2AF2 recognises RNA sequences and how this is modulated by trans-acting RBPs. We measured U2AF2 affinities at hundreds of binding sites, and compared *in vitro* and *in vivo* binding landscapes by mathematical modelling. We found that trans-acting RBPs extensively regulate U2AF2 binding *in vivo*, including enhanced recruitment to 3' splice sites and clearance of introns. Using machine learning, we identified and experimentally validated novel trans-acting RBPs (including FUBP1, CELF6 and PCBP1) that modulate U2AF2 binding and affect splicing outcomes. Our study offers a blueprint for the high-throughput characterisation of *in vitro* mRNP assembly and *in vivo* splicing regulation.

Figure 1. The high-throughput mutagenesis screen provides quantitative splicing information across the RON minigene. Mutagenic PCR creates a library of mutated minigene variants that upon transfection into human cell lines give rise to alternatively spliced transcripts. A unique 15-nt barcode sequence in each minigene (coloured nucleotides) links mutated minigenes and corresponding splicing products, which are characterised by next-generation DNA and RNA sequencing, respectively. A mathematical model allows for a high-throughput identification of splicing-effective mutations.

FUTURE DIRECTIONS

Quantitative description of protein-RNA interactions with the *in vivo* and *in vitro* iCLIP technologies offers an attractive system for understanding the forces of competition and synergy that govern RNP complexes. We focus our efforts on elucidating 3' splice-site definition, which was previously identified as a hotspot for cancer-associated mutations. To achieve this, we are combining *in vivo* and biochemical approaches on a genome-wide scale,

which will yield a systemic understanding of RNP function in splicing regulation.

Reaching into translational research, we are extending our high-throughput mutagenesis approach to study splicing-mediated resistance in paediatric cancer therapy. The knowledge of splicing-effective mutations will enable us to develop prognostic biomarkers and may ultimately lead to new therapy strategies.

SELECTED PUBLICATIONS

Sutandy FXR*, Ebersberger S*, Huang L*, Busch A, Bach M, Kang HS, Fallmann J, Maticzka D, Backofen R, Stadler PF, Zarnack K, Sattler M, Legewie S* and König J. (2018). *In vitro* iCLIP-based modeling uncovers how the splicing factor U2AF2 relies on regulation by cofactors. *Genome Res*, 28: 699–713.

Braun S*, Enculescu M*, Setty ST*, Cortés-López M, de Almeida BP, Sutandy FXR, Schulz L, Busch A, Seiler M, Ebersberger S, Barbosa-Morais NL, Legewie S*, König J* and Zarnack K#. (2018). Decoding a cancer-relevant splicing decision in the RON proto-oncogene using high-throughput mutagenesis. *Nat Commun*, 9: 3315.

Tajnik M, Vigilante A, Braun S, Hänel H, Luscombe NM, Ule J, Zarnack K# and König J#. (2015). Intergenic Alu exonisation facilitates the evolution of tissue-specific transcript ends. *Nucleic Acids Res*, 43: 10492–10505.

* indicates joint contribution
indicates joint correspondence

STEFAN LEGEWIE

“MODELS HELP US TO DISSECT THE COMPLEX CIS-REGULATORY LANDSCAPE OF ALTERNATIVE SPLICING”



EDUCATION

- 2008** PhD in Biophysics, Humboldt University, Berlin
2004 Diploma in Biochemistry, University of Witten/Herdecke

POSITIONS HELD

- Since 2010** Group Leader, Institute of Molecular Biology (IMB), Mainz
2009 – 2010 Group Leader, German Cancer Research Center (DKFZ), Heidelberg
2008 – 2009 Postdoc, Institute for Theoretical Biology, Humboldt University Berlin

GROUP MEMBERS

- Alex Anyaegbunam** PhD Student; since 08/2016
Kolja Becker PhD Student; 08/2013 – 07/2018
Mihaela Enculescu Postdoc; since 10/2013
Rishi Horn Student Assistant; since 10/2018
Sofya Lipnitskaya PhD Student; since 08/2017
Christina Meyer Student Assistant; since 10/2018
Lorenz Ripka PhD Student; since 05/2017

OVERVIEW

Eukaryotic cells sense and process information in order to respond to environmental changes. While the signalling pathways relaying information from the membrane to the nucleus are well characterised, much less is known about decision making at the level of gene expression responses. One focus of our group is to derive a systems-level understanding of gene regulation, which describes: (i) the interplay of signalling pathways and transcription factors in complex gene-regulatory networks; (ii) how the gene expression is co-ordinately controlled at the transcriptional and post-transcriptional levels. We tackle these questions by integrating systematic perturbation screens and multi-OMICS data to derive predictive mathematical models.

A second focus of our group is the quantitative description of cellular heterogeneity. Even genetically identical cells frequently respond in different ways to the same external stimulus, leading to differences in differentiation programs, drug resistance and viral pathogenesis. Together with experimental partners, we employ live-cell imaging approaches to calibrate stochastic and deterministic models of cell population heterogeneity. We employ these models to: (i) derive experimentally testable hypotheses about the causes and consequences of cellular heterogeneity; (ii) better understand therapeutic intervention strategies.

RESEARCH HIGHLIGHTS

Single-cell dynamics of signalling and transcription

We investigated cell-to-cell variability in TGF β /SMAD signalling, which plays a key role in tumorigenesis and metastasis. Together with Alexander Löwer (Darmstadt University), we monitored SMAD2 nuclear translocation in thousands of living MCF10A cells and explained heterogeneous signalling using a quantitative modelling approach, thereby gaining insights into why only a subset of cells induce a migratory response. Along similar

lines, using quantitative imaging and stochastic modelling, we characterised the heterogeneous growth of breast cancer cells by studying the stochastic transcription of a key regulator of estrogen-induced cell proliferation. We found that conventional therapeutic inhibitors of estrogen signalling have limited efficacy, as they induce pronounced gene expression heterogeneity at the single-cell level. This heterogeneity can be overcome by the co-application of small-molecule inhibitors of epigenetic processes. Hence, therapeutic success in cancer treatment could be greater when estrogen receptor antagonists would be applied in combination with other inhibitors.

Reconstruction of alternative splicing networks

Alternative splicing increases protein diversity in eukaryotic cells thereby playing an important role in development and tissue identity but also in diseases such as cancer. Splicing reactions are catalysed by the spliceosome and modulated by auxiliary RNA-binding proteins (RBPs). These guide spliceosome activity, thereby promoting splicing specificity. The RBP U2AF2 is central to alternative splicing decisions as it recruits the spliceosomal machinery to 3' splice sites. To study how U2AF2 recognises RNA sequences, we, together with the König group, established "in vitro iCLIP" experiments in which recombinant RBPs are incubated with long transcripts. Based on titration experiments, we determined U2AF2 affinities at hundreds of binding sites and compared the *in vitro* and *in vivo* binding landscapes by

mathematical modelling. Surprisingly, purified U2AF65 does not exhibit any intrinsic binding preference for 3' splice-sites, suggesting extensive modulation of U2AF2 binding by trans-acting RBPs in living cells. Using machine learning, we identified RNA-binding motifs that are predictive for enhanced or reduced U2AF2 binding and experimentally validated novel trans-acting RBPs controlling U2AF2 binding and splicing outcomes. Our study provided insights into the specificity of alternative splicing decisions.

Mutations causing aberrant splicing are frequently implicated in human diseases including cancer. Together with the König and Zarnack groups, we established a high-throughput screen of randomly mutated minigenes to decode the cis-regulatory landscape that determines alternative splicing of exon 11 in the proto-oncogene MST1R (RON). A combination of kinetic and linear regression modelling enabled us to identify the effects of individual point mutations and to understand how these individual mutations interact to control splicing outcomes. In fact, we identified more than 1,000 mutations affecting RON exon 11 skipping, which corresponds to the pathological isoform RON Δ 165 (Figure 1). Importantly, the effects correlate with RON alternative splicing in cancer patients bearing the same mutations. Our results, therefore, offer insights into splicing regulation and the impact of mutations on alternative splicing in cancer. In the future, we plan to further characterise trans-acting RBPs that recognise cis-regulatory elements defined by the mutagenesis screen.




Figure 1. Regression modelling determines more than 1,900 single mutation effects in the MST1R (RON) proto-oncogene. Experimental data (left): Using RNA-Seq, splice isoforms were quantified for ~5,000 minigene variants, each harbouring a unique set of combined point mutations. A kinetic model of splicing reactions (middle) reveals that point mutation effects show additive behaviour if the data is properly normalised. Thus, a linear regression model (right) was formulated, in which the measured effect of combined mutations equals the sum of single mutation effects. The single mutation effects are inferred by fitting this model to the data. See Braun et al (2018) for details.

FUTURE DIRECTIONS

We plan to further refine and develop existing models of signalling and gene expression. For instance, we are investigating the stochastic dynamics of TGF β /SMAD signalling and studying additional prototypical splicing events. With this, we hope to convert these general insights into a splice code, which describes splicing

outcomes based on sequence information and RBP expression patterns. Moreover, we have begun to model the dynamics of DNA (de)methylation to better understand how this important epigenetic mark can be set and erased to tune gene activity.

SELECTED PUBLICATIONS

Fritsch C*, Baumgärtner S*, Kuban M, Steinshorn D, Reid G[#] and Legewie S*. (2018). Estrogen-dependent control and cell-to-cell variability of transcriptional bursting. *Mol Syst Biol*, 14: e7678.

* indicates joint contribution
[#] indicates joint correspondence

Braun S*, Enculescu M*, Setty ST*, Cortés-López M, de Almeida BP, Sutandy FXR, Schulz L, Busch A, Seiler M, Ebersberger S, Barbosa-Morais NL, Legewie S*, König J[#] and Zarnack K[#]. (2018). Decoding a cancer-relevant splicing decision in the RON proto-oncogene using high-throughput mutagenesis. *Nat Commun*, 9: 3315.

Kamenz J, Mihaljev T, Kubis A, Legewie S[#] and Hauf S*. (2015). Robust ordering of anaphase events by adaptive thresholds and competing degradation pathways. *Mol Cell*, 60: 446–459.

EDWARD LEMKE

“DISORDERED PROTEINS
ENCODE SPECIFICITY
AND SELECTIVITY INTO
MOLECULAR
MECHANISMS”



EDUCATION

- 2005** PhD, Max Planck Institute for Biophysical Chemistry & University of Göttingen
- 2001** Diploma in Chemistry, Technical University of Berlin
- 2001** MSc in Biochemistry, University of Oklahoma

POSITIONS HELD

- Since 2018** Adjunct Director, Institute of Molecular Biology (IMB), Mainz
Professor of Synthetic Biophysics, Johannes Gutenberg University Mainz
- Since 2009** Group Leader, European Molecular Biology Laboratory (EMBL), Heidelberg
- 2005 – 2008** Research Associate (Postdoc), The Scripps Research Institute, La Jolla

GROUP MEMBERS

- Joana Caria** Lab Manager; since 04/2017
- Giorgia Celetti** Technology Engineer; since 06/2016
- Christine Koehler** PhD Student; since 01/2017
- Sofya Mikhaleva** PhD Student; since 09/2015
- Giulia Paci** PhD Student; since 10/2014
- Panagiotis Patsis** PhD Student; since 11/2018
- Christopher Reinkemeier** PhD Student; since 09/2016
- Piau Siong Tan** Postdoc; since 04/2015
- Miao Yu** Postdoc; since 10/2018

OVERVIEW

We focus on studying intrinsically disordered proteins (IDPs), which constitute up to 50% of the eukaryotic proteome. IDPs are found in many vital biological processes, such as nucleocytoplasmic transport, transcription and gene regulation. The ability of IDPs to exist in multiple conformations is considered a major driving force behind their enrichment during evolution in eukaryotes. Studying biological machineries containing such dynamic proteins is a major hurdle for conventional technologies. Because of this and as they are hard to visualise, IDPs are termed the dark proteome. Using a question-driven, multidisciplinary approach paired with novel tool development, we have made major strides in understanding the biological dynamics of such systems from the single molecule to the whole cell level. Fluorescence tools are ideally suited to study the plasticity of IDPs, since their non-invasive character permits smooth transition between *in vitro* (biochemical) and *in vivo* (in cell) studies. In particular, single molecule and super-resolution techniques are powerful tools for studying the spatial and temporal heterogeneities that are intrinsic to complex biological systems. We synergistically combine this effort with advanced tool developments in chemical biology, microfluidics and microscope engineering to increase the throughput, strength and sensitivity of the approach as a whole.

RESEARCH HIGHLIGHTS

Our strong focus on the mechanistic understanding of IDPs using single molecule and super-resolution tools is both driven and driving novel tool developments for “in-cell/*in situ* structural biology”. This comprises of a synergistic effort of chemical/synthetic biology and precision fluorescence-based technology/nanoscopy/microfluidics development (Figure 1).

A major technical breakthrough of my lab was the ability to engineer “click”-able functionalities into any protein both *in vitro* and *in vivo*. This genetic code expansion (GCE) approach has the potential to become a true GFP (fusion protein) surrogate strategy, with the major advantages being that superior synthetic dyes can be coupled with residue-specific precision anywhere in a protein. This opens up new avenues in single-molecule fluorescence and super-resolution microscopy.

These precision tools enable us to make even the most complex molecular machinery visible to our core methodologies, which are based on time-resolved multiparameter and nanoscopy tools. This enables innovative approaches to study the heterogeneity of IDPs in biology. More recently, we discovered a distinct ultrafast protein-protein interaction mechanism that can explain how nuclear pore complexes (NPCs) can efficiently fulfil their central role in cellular logistics and how nuclear transport can be both fast and selective at the same time. We also determined how this function can coexist with other nuclear transport mechanisms that provide a platform for cargo undocking. These findings provided a leap forward in our understanding of how IDPs maintain different functionalities through conformational changes despite the normal requirement for rigid molecular specificity.

Despite our advancing technologies for *in situ* science, we always consider it important to perform studies on reconstituted systems (*in vitro*/biochemical) to understand our biological problems in a bottom-up fashion and

complement our results from our *in situ* studies. To achieve this, we worked on various aspects e.g. i) utilising microfluidics to integrate Lab-on-chip technology into our workflows and ii) developing a versatile baculovirus-based platform, which combines the benefits from GCE technology and the versatility of click chemistry with the strength of recombinant protein engineering. This technology now enables *in vitro*/bottom up/reconstitution based biological project design that was previously unachievable *in situ*. Our work is accompanied by rigorous analysis on the physicochemical properties of IDPs and examines to what extent simple, known polymer concepts, such as phase separation, can be used to describe the function of such biopolymers *in vivo*.




Figure 1. Schematic representation of technologies used in the Lemke Lab to study the Dark Proteome.

FUTURE DIRECTIONS

IDPs are highly multifunctional. Due to their multivalency and large disordered regions they can function as dynamic scaffold platforms. We combine chemical and synthetic biology approaches to enable non-invasive, multi-color high- and super-resolution studies of activity-dependent changes of protein conformation in living cells, enabling fluorescence driven *in situ* structural biology. The key point is that the enhanced spatial and temporal resolution offered by our fluorescent methods will enable us to detect rare events and unexpected behaviours inside cells. We want to

use this to identify and understand IDP multifunctionalities that are clearly distinguishable from their normal mode of action, as for example nucleoporins (Nups) in the nuclear pore complex (NPC). In fact, many IDP-Nups have roles in pathogen-host interactions and have been suggested to shuttle away from the NPC to function in gene regulatory processes. A prominent example of IDP-Nup action distinct from its normal mode in the NPC is Nup98, since several genetic fusions of Nup98 with transcription factors are related to leukaemia.

SELECTED PUBLICATIONS

Tan PS, Aramburu IV, Mercadante D, Tyagi S, Chowdhury A, Spitz D, Shammass SL, Gräter F and Lemke EA. (2018). Two differential binding mechanisms of FG-nucleoporins and nuclear transport receptors. *Cell Rep*, 22: 3660–3671.

Nikić I, Estrada Girona G, Kang JH, Paci G, Mikhaleva S, Koehler C, Shymanska NV, Ventura Santos C, Spitz D and Lemke EA. (2016). Debugging eukaryotic genetic code expansion for site-specific click-PAINT super-resolution microscopy. *Angew Chemie Int Ed*, 55: 16172–16176.

Milles S, Mercadante D, Aramburu IV, Jensen MR, Banterle N, Koehler C, Tyagi S, Clarke J, Shammass SL, Blackledge M, Gräter F and Lemke EA. (2015). Plasticity of an ultrafast interaction between nucleoporins and nuclear transport receptors. *Cell*, 163: 734–745.

BRIAN LUKE

“WE ARE ONLY BEGINNING TO UNDERSTAND THE REGULATION OF RNA IN A CHROMATIN ENVIRONMENT”



EDUCATION

- 2005** PhD in Biochemistry, Swiss Federal Institute of Technology Zurich (ETH)
1999 BSc in Biology, Queen's University, Ontario

POSITIONS HELD

- Since 2017** Adjunct Director, Institute of Molecular Biology (IMB), Mainz
 Heisenberg Professor, Johannes Gutenberg University Mainz
2014 – 2017 Group Leader, Institute of Molecular Biology (IMB), Mainz
2009 – 2014 Group Leader, Centre for Molecular Biology (ZMBH), University of Heidelberg
2005 – 2009 Postdoc, Swiss Federal Institute of Technology Lausanne (EPFL)
2005 Postdoc, Swiss Federal Institute of Technology Zurich (ETH)

GROUP MEMBERS

- Katharina Bender** PhD Student; 03/2015 – 10/2018
Fabio Bento PhD Student; since 04/2017
Stefanie Reimann Technician; since 01/2015
Vanessa Kellner PhD Student; 03/2015 – 05/2018
Arianna Lockhart Postdoc; since 04/2018
Sarah Luke-Glaser Staff Scientist; 10/2016 – 12/2018
Stefano Misino PhD Student; since 10/2016
Vanessa Pires PhD Student; since 08/2017
Natalie Schindler Postdoc; since 12/2017
Simone Snead Lab Manager; 04/2017 – 10/2018
Christiane Stürzbecher Personal Assistant; since 04/2017
Matthias Tonn PhD Student; since 04/2017
Olga Vydzhak PhD Student; since 02/2016
Tina Wagner PhD Student; since 11/2015

OVERVIEW

It has become increasingly clear that RNA plays a critical role in a myriad of DNA transactions ranging from transcription regulation to DNA repair and chromosome 3D structure. RNA has the ability to form hybrid molecules with DNA in the form of R-loops, RNA-DNA triplexes and ribonucleoside monophosphate (rNMP) insertions directly into the DNA strands. We are only beginning to understand how these RNA moieties are regulated as well as the consequences that arise when they are misregulated. We are currently focused on understanding how the RNase H enzymes are coordinated in RNA-DNA hybrid removal in terms of both time and space. Currently, there is very little understanding regarding the regulation of these critical RNA-DNA hybrid regulatory enzymes. In addition to the regulation of the RNase H enzymes, we are trying to understand how the replication machinery reacts when RNA-DNA hybrids are not adequately eliminated. In this respect, a recently described RNase H2 separation of function allele is allowing us to interrogate the differences between rNMPs and R-loops. We use a variety of biochemical, proteomic and genetic techniques to address these mechanistic questions. The recent implication of RNase H mutations in human disease makes these studies pertinent to eventual medical applications.



RESEARCH HIGHLIGHTS

We have had a longstanding interest in understanding how the ubiquitin ligase, Rtt101^{Mms1} contributes to genome stability and DNA replication fidelity. We have demonstrated that the conserved Rtt101 ligase becomes essential when rNMPs accumulate (Figure 1A). This genetic interaction is relevant, as mutations in RNase H2, which removes rNMPs from the genome, have been implicated in neurological disorders as well as human cancer. We used a mass spectrometry-coupled ubiquitin remnant profiling approach to identify the relevant target(s) of Rtt101 when

rNMPs accumulate. We identified and confirmed that the DNA polymerase epsilon subunit, Dpb2, gets ubiquitylated in a Rtt101 dependent manner when ribonucleotides accumulate (Figure 1B). Follow-up biochemical fractionations have demonstrated that Rtt101 promotes the removal of Dpb2, as well as other DNA polymerase epsilon subunits, from chromatin. Based on these results, we propose a hypothesis whereby DNA damage-induced replication stress on the leading strand triggers the eviction of DNA polymerase epsilon in a Rtt101 ubiquitylation dependent manner. Based on our previous studies, we speculate that polymerase removal is essential to promote homologous recombination-mediated restart of stalled replication forks.

RNase H enzymes are dedicated to the nucleolytic removal of RNA that is hybridised to DNA. RNase H1 is a monomeric enzyme that degrades RNAs hybridised to DNA with the involvement of 4 or more base pairs, and is largely limited to R-loop metabolism. RNaseH2, on the other hand, is a heterotrimeric enzyme that can eliminate both R-loops and rNMPs. By creating cell cycle-regulated alleles of the RNase H enzymes we have been able to determine that RNase H1 removes R-loops in the S phase of the cell cycle. RNase H2, however, functions in the G2 phase of the cell cycle where it carries out ribonucleotide excision repair as well as R-loop removal. These results suggest that RNase H2 may also promote a post-replicative repair function.

Figure 1. A) When the catalytic subunit of RNase H2 (Rnh201) is degraded in the presence of auxin (IAA), and rNMPs accumulate, Rtt101 becomes essential (see bottom row). B) In the absence of Rtt101, Dpb2 is under-ubiquitylated in a genetic background with high rNMP levels as assessed by quantitative mass spectrometry and ubiquitin profiling.



Telomeres are transcribed into a non-coding RNA, referred to as TERRA (Telomeric repeat-containing RNA). We have demonstrated that TERRA R-loops accumulate at short telomeres and promote homology-directed repair due to the inability of RNase H2 to localise to those telomeres. We have used a proteomics approach to assess which proteins may localise to short telomeres in an RNA dependent manner. We identified Npl3, an hnRNP-like protein, as a novel telomere binding factor. Npl3 is specifically recruited to short telomeres in an R-loop dependent manner. In the absence of Npl3, cells enter replicative senescence with accelerated kinetics, suggesting the recruitment of Npl3 to short telomeres is essential to regulate senescence onset by directly regulating telomeres. Our preliminary data indicate that Npl3 may be promoting the formation of R-loops, which in turn drives telomeric recombination.

Finally, we have shown that when repair defective yeast cells are exposed to DNA damaging agents, they undergo checkpoint adaptation. The adapted cells acquire drug resistance and are aneuploid. We have now shown that the combination of genotoxic agents together with targeting of adaptation and aneuploidy, leads to a synergistic cytotoxicity specifically of repair defective cells. These results are highly relevant for cancer chemotherapy.

FUTURE DIRECTIONS

The function of Dpb2 ubiquitylation will be investigated with regards to how it affects DNA replication and repair dynamics by creating non-ubiquitylatable Dpb2 alleles and analysing replication and repair. We will also try to determine whether the ubiquitylation of Dpb2 leads to degradation of the protein. Finally, we will investigate upstream signalling events that may trigger the modification of Dpb2.

The genetic data that we have generated with the RNase H cell cycle alleles will be substantiated with biochemical experi-

ments to determine levels of R-loops and rNMPs in the genome when the alleles are expressed. Using a combination of genetic experiments and imaging we will test the potential post-replicative repair function of RNase H2. We will use the telomere as a model locus to elucidate how R-loops drive recombination. We will test the hypothesis that for recombination at telomeres an R-loop must be present but then eventually removed. Npl3 and Rnh1 are candidates to examine in this context.

SELECTED PUBLICATIONS

Graf M*, Bonetti D*, Lockhart A*, Serhal K, Kellner V, Maicher A, Jolivet P, Teixeira MT and Luke B. (2017). Telomere length determines TERRA and R-loop regulation through the cell cycle. *Cell*, 170: 72–85.

* indicates joint contribution
indicates joint correspondence

Lafuente-Barquero J, Luke-Glaser S, Graf M, Silva S, Gómez-González B, Lockhart A, Lisby M, Aguilera A and Luke B. (2017). The Smc5/6 complex regulates the yeast Mph1 helicase at RNA-DNA hybrid-mediated DNA damage. *PLOS Genet*, 13: e1007136.

Buser R*, Kellner V*, Melnik A, Wilson-Zbinden C, Schellhaas R, Kastner L, Piwko W, Dees M, Picotti P, Maric M, Labib K, Luke B* and Peter M#. (2016). The replisome-coupled E3 ubiquitin ligase Rtt101^{Mms22} counteracts Mrc1 function to tolerate genotoxic stress. *PLOS Genet*, 12: e1005843.

CHRISTOF NIEHRS

“WE ARE A STEP CLOSER TO UNDERSTANDING HOW R-LOOPS ACT AS EPIGENETIC GENE REGULATORS”



EDUCATION

- 1997** Habilitation in Biology, University of Heidelberg
- 1990** PhD in Biology, European Molecular Biology Laboratory (EMBL) & University of Heidelberg
- 1985** Diploma in Biochemistry, Free University of Berlin

POSITIONS HELD

- Since 2010** Founding & Scientific Director, Institute of Molecular Biology (IMB), Mainz
Professor, Johannes Gutenberg University Mainz
- 2010 – 2015** Executive Director, Institute of "Molecular Embryology" (IMB), Mainz
- Since 2000** Professor of Molecular Embryology, German Cancer Research Center (DKFZ), Heidelberg
- Since 1994** Head of Division "Molecular Embryology", German Cancer Research Center (DKFZ), Heidelberg
- 1990 – 1993** Postdoc, University of California Los Angeles (UCLA)

GROUP MEMBERS

- Khelifa Arab** Postdoc; since 11/2011
- Amitava Basu** Postdoc; since 03/2018
- Anne Baumgärtner** PhD Student; since 12/2016
- Tamara Dehn** Animal Caretaker; since 06/2011
- Anna Luise Ernst** PhD Student; since 04/2015
- David Fournier** Postdoc; since 09/2018
- Dandan Han** PhD Student; since 12/2013
- Laura Krebs** Technician; since 09/2015
- Medhavi Mallick** PhD Student; 08/2012 – 09/2018
- Carmen Meyer** Technician; since 05/2018
- Michael Musheev** Postdoc; since 07/2011
- Regina Otto** Personal Assistant; since 09/2015
- Eleftheria Parasyraki** PhD Student; since 09/2018
- Mihika Pradhan** PhD Student; since 12/2016
- Sandra Rölle** Lab Manager; since 04/2011
- Andrea Schäfer** Postdoc; since 09/2010
- Carola Scholz** Technician; since 05/2015
- Lars Schomacher** Postdoc; since 07/2011
- Katrin Schüle** PhD Student; since 01/2016
- Philipp Trnka** PhD Student; since 11/ 2016
- Viviana Vastolo** Postdoc; since 09/2015

OVERVIEW

In the genome of many multicellular organisms, DNA methylation is a common epigenetic mark associated with gene silencing. DNA methylation is a dynamic process and can be reversed by enzymatic demethylation, a process that is still incompletely understood. DNA demethylation is a widespread phenomenon occurring in plants as well as in animals during development, in the adult and during the somatic cell reprogramming of pluripotency genes. We showed that Growth arrest and DNA damage 45A (GADD45A) is a key player in active DNA demethylation and acts via DNA repair. One goal of our research is to analyse the mechanism of DNA demethylation as well as the role played by GADD45A in development. Our results indicate that GADD45A acts as an adapter protein, which directs DNA methylation machinery to specific loci. To address GADD45A function, we use biochemical, molecular biology and cell biological approaches, employing the mouse and frog model systems as well as embryonic stem cells.

RESEARCH HIGHLIGHTS

DNA methylation at 5-methylcytosine (5mC) of CpGs plays important roles in regulating gene expression, genomic imprinting, X-chromosome inactivation, genomic instability, embryonic development, and cancer. DNA methylation is reversible by enzymatic active DNA demethylation, with examples in plants, animal development, cancer, and immune cells. Yet, the molecular mechanisms underlying active demethylation are only beginning to be understood.

We have shown previously that the stress response protein *Gadd45a* mediates active DNA demethylation. GADD45 proteins are multifunctional and regulate a range of cellular processes, including DNA repair, proliferation, apoptosis, and differentiation. GADD45A directly interacts with the two key enzymes of the DNA

demethylation machinery, TET (Ten-eleven translocation) and TDG (thymine-DNA glycosylase), to enhance turnover of oxidized cytosines. Therefore, GADD45A acts as an adapter, which recruits DNA modifying enzymes to specific sites in the genome and promotes local demethylation.

To be directed to specific genomic loci, GADD45A relies on bridging factors, which include RNA. We previously showed that expression of the tumour suppressor *TCF21* is activated by a lncRNA, termed TARID (for *TCF21* antisense RNA inducing demethylation), which is transcribed in antisense orientation to *TCF21*. TARID recruits the DNA demethylation machinery to the *TCF21* promoter, leading to decreased DNA methylation and increased expression of *TCF21*. TARID-dependent activation of *TCF21* transcription is brought about by recruitment of GADD45A, which tethers TDG and TET proteins to specific genomic sites to direct base excision repair mediated DNA demethylation.

How does TARID interact with the *TCF21* promoter? We hypothesised that being an antisense transcript, TARID could form an R-loop at a CpG island (CGI) in the promoter of *TCF21*, which is recognised by GADD45A. R-loops are naturally occurring three-stranded DNA:RNA hybrids and have recently gained increasing attention, since they play important roles in transcription, RNA processing, DNA damage, and genome stability. Genome-wide mapping showed that they exist throughout the genome under physiological conditions in diverse organisms. R-loops can be detrimental to cells because they can induce DNA damage and genome instability and have been associated with neurodegenerative disease. Conversely, R-loops also play a regulatory role in various

nuclear processes, notably in the regulation of gene expression.

We now found that TARID indeed forms an R-loop at the *TCF21* promoter, which is bound by GADD45A to trigger local DNA demethylation and *TCF21* expression. TARID transcription, R-loop formation, DNA demethylation, and *TCF21* expression proceed sequentially during the cell cycle to allow both the sense and antisense RNAs to be expressed without hindrance. In support of a generic action of GADD45A we found that oxidized DNA demethylation intermediates are enriched at genomic R-loops and that their levels increase upon RNase H1 depletion. Genomic profiling in embryonic stem cells identified thousands of R-loop-dependent TET1 binding sites at CGIs. Our study supports to notion, that R-loops associated with certain classes of RNAs can act as *bona fide* epigenetic regulators. Conceptually, R-loops are attractive candidates because analogous to bacterial guide RNAs, which form R-loops to direct CAS9 endonuclease, mammalian R-loops may direct epigenetic modifiers in a sequence-specific manner via base pairing to genomic loci. Our study indicates that GADD45A is an epigenetic reader of such regulatory R-loops at CGIs (Figure 1).




Figure 1. An R-loop formed between an antisense lncRNA and a cognate coding gene assembles an RNP complex including an epigenetic reader (GADD45A) and writer (TET DNA demethylase). Demethylated CpGs (lollipops) at a CpG island (CGI) are indicated.

FUTURE DIRECTIONS

Our discovery of GADD45A as an R-loop reader raises new questions. How are R-loops decoded in embryonic stem cell pluripotency and differentiation? Which R-loop regions in ESCs are engaged in epigenetic regulation and what are their molecular determinants? To which R-loops is TET1 guided? How are regula-

tory R-loops erased? Are there other specific R-loop readers and what are their functions? We will approach these questions in embryonic stem cells where R-loops have been well documented and which are easily amenable to genetic experimentation.

SELECTED PUBLICATIONS

Arab K, Karaulanov E, Musheev M, Trnka P, Schäfer A, Grummt I and Niehrs C. (2019). GADD45A binds R-loops and recruits TET1 to CpG island promoters. *Nat Genet*, doi: 10.1038/s41588-018-0306-6.

Schäfer A, Mekker B, Mallick M, Vastolo V, Karaulanov E, Sebastian D, von der Lippen C, Epe B, Downes DJ, Scholz C and Niehrs C. (2018). Impaired DNA demethylation of C/EBP sites causes premature aging. *Genes Dev*, 32: 742–762.

Arab K, Park YJ, Lindroth AM, Schäfer A, Oakes C, Weichenhan D, Lukanova A, Lundin E, Risch A, Meister M, Dienemann H, Dyckhoff G, Herold-Mende C, Grummt I[#], Niehrs C[#] and Plass C[#]. (2014). Long noncoding RNA TARID directs demethylation and activation of the tumor suppressor *TCF21* via GADD45A. *Mol Cell*, 55: 604–614.

[#] indicates joint correspondence

HOLGER RICHLY

“WE INVESTIGATE THE
IMPACT OF DNA REPAIR
MECHANISMS IN DISEASE
AND ORGANISMAL
AGEING”



EDUCATION

- 2005** PhD in Biology, Ludwig Maximilian University (LMU), Munich
- 2000** Diploma in Biochemistry, Ruhr University, Bochum

POSITIONS HELD

- Since 2011** Group Leader, Institute of Molecular Biology (IMB), Mainz
- 2005 – 2011** Postdoc, Centre for Genomic Regulation (CRG), Barcelona

GROUP MEMBERS

- Daniel Henning** Master's Student; 04/2018 – 11/2018
- Aysegül Kaymak** PhD Student; 02/2013 – 04/2018
- Rebeca Medina** Technician; 08/2013 – 11/2018
- Jens Stadler** PhD Student; since 09/2014

OVERVIEW

The research undertaken in my laboratory aims at deciphering molecular pathways that underlie chromatin signalling networks that regulate physiological processes such as DNA repair and organismal ageing. Our scientific approach relies largely on dissecting the functions of diverse chromatin components, as for example epigenetic players, biochemically in cell culture systems and by applying high resolution microscopy. We complement our *in vitro* approach by employing genetics and RNAi screening techniques in *Caenorhabditis elegans*.

RESEARCH HIGHLIGHTS

Epigenetic networks govern most cellular processes that take place in a chromatin environment, for example differentiation, DNA repair and replication. Our research provides evidence for how epigenetic factors act in concert with DNA repair factors. In our investigations studying DNA repair, we have largely concentrated on one particular histone mark, the mono-ubiquitylation of histone H2A at lysine 119 (H2A-ubiquitin). H2A-ubiquitylation is a hallmark of signalling cascades as part of the DNA damage response. We have demonstrated that timing of DNA repair specific E3 ligases is an important feature of nucleotide excision repair (NER) and we have discussed a new concept of remodelling E3 ligase complexes at chromatin during DNA lesion recognition. In brief, we discovered that H2A-ubiquitin is catalysed predominantly by a novel E3 ligase complex (UV-RING1B complex) that operates early during lesion recognition (Figure 1). ZRF1 tethers to the H2A-ubiquitin mark at the damage site and mediates the remodelling of the UV-RING1B complex, a process that we have coined *on-site remodelling*.

We have further demonstrated that on-site remodelling in the global genomic branch of NER is confined to specific nuclear

regions. More recently, we have shown that ZRF1, apart from remodelling multi-protein complexes, is also engaged in the decondensation of chromatin. ZRF1 recruits the endoribonuclease DICER to the DNA damage sites and both proteins in conjunction with PARP1 facilitate the remodelling of chromatin (Figure 1). We have further demonstrated that DICER recruits the methyltransferase MMSET to the DNA damage site, which catalyzes the dimethylation of histone 4 at lysine20 (H4K20me2). This chromatin mark tethers the DNA repair factor XPA via the adaptor proteins 53BP1 and RPA2 (Figure 1). More recently, we have examined the concerted function of ubiquitylation events and DNA incision at the DNA damage site.

Furthermore, we are interested in understanding the impact of gene regulation and DNA repair mechanisms during organismal ageing. To this end we investigate aging in the nematode *C. elegans*, employing sophisticated RNAi screening techniques, genetics and high resolution microscopy. We performed the first late-life RNAi screen designed to discover novel longevity genes that exhibit antagonistic pleiotropy in *C. elegans*. We identified that post-reproductive inactivation of genes required for autophagosome nucleation, such as the *Atg6/VPS30/beclin-1* ortholog *bec-1*, led to a strong lifespan increase of up to 60% post-RNAi initiation. We could further show that while the process of vesicle nucleation is still active and possibly enhanced in old worms, the process is blocked downstream of autophagosome biogenesis at the step of autolysosomal degradation. Moreover, we found that post-reproductive inactivation of auto-

phagosome nucleation extends lifespan primarily through the neurons. Currently we are exploring the functions of DNA repair factors in organismal ageing. To this end we have carried out a late-life RNAi screening of factors from different DNA repair systems. We are now investigating candidate DNA repair factors in detail to reveal their significance in the ageing process.




Figure 1. Spatiotemporal regulation of chromatin factors and ubiquitylation events during NER. (top) The assembly of the UV-RING1B complex at DNA damage sites causes mono-ubiquitylation of histone H2A. ZRF1 is recruited to the damage site via XPC and tethers to H2A-ubiquitin causing on-site remodelling of the UV-RING1B E3 ligase complex. The newly-established DDB-CUL4A complex catalyzes the polyubiquitylation of various substrates and most importantly XPC, which is thereby stabilised at the damage site. (middle) ZRF1 recruits DICER to DNA damage sites and decondenses chromatin in conjunction with PARP1. (bottom) DICER recruits MMSET to catalyze H4K20 methylation at the DNA damage site. H4K20me2 tethers XPA via 53BP1 and RPA2.

FUTURE DIRECTIONS

In the future, we will further prioritise the research on DNA repair in the NER pathway and the investigation of organismal ageing. One of our main aims is to understand the chromatin signalling network underlying DNA damage recognition in the NER pathway and the transition from recognition to the verification of DNA damage. In particular, we will analyse how ubiquitin signalling cascades crosstalk to other chromatin factors

and histone marks. Furthermore, we will investigate the function of K63-linked polyubiquitylation in NER, which presumably provides a means of recruiting repair factors. To extend our studies in organismal ageing, we plan to study how chromatin factors and environmental cues extend the lifespan and the health span of *C. elegans*. In particular, we will examine late life functions of metabolic genes.

SELECTED PUBLICATIONS

Chitale S and Richly H. (2018). DICER- and MMSET-catalyzed H4K20me2 recruits the nucleotide excision repair factor XPA to DNA damage sites. *J Cell Biol*, 217: 527–540.

* indicates joint contribution

Wilhelm T*, Byrne J*, Medina R, Kolundžić E, Geisinger J, Hajduskova M, Tursun B and Richly H. (2017). Neuronal inhibition of the autophagy nucleation complex extends life span in post-reproductive *C. elegans*. *Genes Dev*, 31: 1561–1572.

Gracheva E, Chitale S, Wilhelm T, Rapp A, Byrne J, Stadler J, Medina R, Cardoso MC and Richly H. (2016). ZRF1 mediates remodeling of E3 ligases at DNA lesion sites during nucleotide excision repair. *J Cell Biol*, 213: 185–200.

JEAN-YVES ROIGNANT

“WE ARE ON THE WAY TO CRACK THE EPITRANSCRIPTOME”



EDUCATION

- 2003** PhD in Developmental Biology, Jacques Monod Institute, Paris
- 1998** MSc in Developmental Biology, Paris VII
- 1997** BSc in Molecular and Cellular Biology, Rennes University

POSITIONS HELD

- Since 2012** Group Leader, Institute of Molecular Biology (IMB), Mainz
- 2008 – 2011** Research Associate, Skirball Institute of Biomolecular Medicine, New York University
- 2003 – 2008** Postdoc, Skirball Institute of Biomolecular Medicine, New York University

GROUP MEMBERS

- Praveen Bawankar** Postdoc; since 05/2018
- Nadja Dinges** PhD Student; 03/2014 – 03/2018
- Annabelle Dold** PhD Student; since 01/2015
- Nadine Körtel** Master's Student; 02/2018 – 10/2018
- Giriram Kumar** PhD Student; since 04/2014
- Jessica Leismann** Master's Student; since 04/2018
- Tina Lence** PhD Student; since 07/2013
- Violeta Morin** Technician; since 05/2015
- Miriam Mulorz** Master's Student; since 07/2018
- Chiara Paolantoni** PhD Student; since 10/2017
- Mariangela Spagnuolo** PhD Student; since 02/2018
- Michael Stock** PhD Student; since 06/2018
- Lina Worpenberg** PhD Student; since 08/2017

OVERVIEW

Our work focuses on elucidating the functions and mechanisms of RNA modifications, a novel layer of post-transcriptional gene regulation also known as epitranscriptomics. Modifications of RNA are very abundant, as more than 170 types have been detected in the last sixty years. However, understanding of their biological significance has been hampered by a lag in the technology needed for their identification. As a result, very little is known about their role in messenger and non-coding RNAs. Over the past few years, work from us and other groups has contributed to deciphering the role of mRNA modifications in gene regulation, revealing their importance in several physiological processes and diseases. Current projects in the lab deal with the m⁶A mRNA modification as well as less characterised modifications. For our research, we use *Drosophila melanogaster* as a model organism and employ state of the art approaches including molecular biology and classical genetics combined with high throughput techniques and computational tools.

RESEARCH HIGHLIGHTS

Elucidation of the roles, mechanisms and targets of the m⁶A mRNA modification

The widespread roles of m⁶A RNA modification in the regulation of post-transcriptional gene expression have recently been brought to light. From yeast to mammals, m⁶A has been shown to regulate pre-mRNA splicing, translation and mRNA decay. However, the precise mechanisms of this modification in these processes still remain to be determined. Furthermore, the physiological functions of m⁶A in multi-cellular organisms have not yet been fully investigated. To address these questions, we have characterised the m⁶A pathway in *Drosophila*. We found that a conserved m⁶A methyltransferase complex controls alternative splicing in both *Drosophila* cells and *in vivo*. As in mammals, components

of the complex are ubiquitously expressed but show significant enrichment in the nervous system, which is consistent with the high level of m⁶A in this tissue. We find that flies mutant for *Mettl3* are viable but suffer from severe locomotion defects due to impaired neuronal functions. A synaptic overgrowth was observed at neuromuscular junctions (Figure 1). Components of the m⁶A methyltransferase complex also control the female-specific splicing of the Sex lethal (*Sxl*) transcript and of its downstream targets, revealing a role for this modification in sex determination and dosage compensation. We have recently identified a new member of the methyltransferase complex that we have named Flacc for Fl(2)d Associated Complex Component. Our work suggests that Flacc serves as an adapter to link Spenito with other components of the methyltransferase complex. We demonstrated that this association is critical for m⁶A deposition. In addition, we have developed a new approach to map m⁶A *in vivo* as current techniques were not adapted for *in vivo* studies. We are currently taking advantage of this approach to characterise m⁶A molecular targets and their functions during organismal development.

Functions of Pseudouridine synthase 7 (*Pus7*) in neuronal behaviour

For decades, pseudouridine has been known to be the most abundant modification on transfer RNA (tRNA) and ribosomal RNA (rRNA) and more recently was found to be also widespread in mRNA molecules. In humans, pseudouridine is catalysed via 13 distinct pseudouridine synthases (PUSs), each of which has distinct specificities, most of them only poorly understood. In collaboration with the groups

of Schraga Schwartz (Weizmann Institute) and Arjan de Brouwer (Radboud University) we have identified several human patients with distinct *Pus7* mutations. All these individuals have intellectual disabilities with speech delay, short stature, microcephaly, and aggressive behaviour. We showed that the disease-related variants lead to abolishment of *PUS7* activity on both tRNA and mRNA substrates. Moreover, we generated a *Pus7* knockout in *Drosophila melanogaster*. Mutant flies are viable but display a number of behavioural defects, including increased activity, disorientation, and aggressiveness supporting the idea that neurological defects are caused by *PUS7* variants. Therefore, our findings demonstrate that RNA pseudouridylation by *PUS7* is essential for proper neuronal development and function. We are currently looking for the relevant targets of *Pus7* in mediating its neuronal defects, both in *Drosophila* and in patient cells.




Figure 1. Lack of m⁶A modification results in axonal growth at neuromuscular junctions. Representative confocal images of muscle – 6/7 NMJ synapses of abdominal hemisegment A2 for the indicated genotypes labelled with anti-Synaptotagmin (green) and HRP (red) to reveal the synaptic vesicles and the neuronal membrane. *Mettl3*^{Δcat}/*Mettl3*^{null} mutants display synaptic outgrowths. Scale bar: 20 μm.

FUTURE DIRECTIONS

Our long-term goal is to decipher the chemical code decorating mRNA and how it impacts gene regulation during development and diseases. We are currently performing CRISPR/Cas9 based-screens to identify novel functions for specific RNA modi-

fications. Our study should expose a new layer of regulation of gene expression that may have important implications for understanding key cellular processes that dictate cell fate.

SELECTED PUBLICATIONS

Knuckles P, Lence T, Haussmann IU, Jacob D, Kreim N, Carl SH, Masiello I, Hares T, Villaseñor R, Hess D, Andrade-Navarro MA, Biggiogera M, Helm M, Soller M, Bühler M and Roignant JY. (2018). Zc3h13/Flacc is required for adenosine methylation by bridging the mRNA-binding factor Rbm15/Spenito to the m⁶A machinery component Wtap/Fl(2)d. *Genes Dev*, 32: 415–429.

Lence T, Akhtar J, Bayer M, Schmid K, Spindler L, Ho CH, Kreim N, Andrade-Navarro MA, Poeck B, Helm M and Roignant JY. (2016). m⁶A modulates neuronal functions and sex determination in *Drosophila*. *Nature*, 540: 242–247.

Malone CD, Mestdagh C, Akhtar J, Kreim N, Deinhard P, Sachidanandam R, Treisman J and Roignant JY. (2014). The exon junction complex controls transposable element activity by ensuring faithful splicing of the *piwi* transcript. *Genes Dev*, 28: 1786–1799.

VASSILIS ROUKOS

“WE USE STATE-OF-THE-ART MICROSCOPY AND GENOMICS TO UNDERSTAND THE GENESIS OF GENOME REARRANGEMENTS”



EDUCATION

- 2008** PhD in Molecular Biology & Cytogenetics, Medical School, University of Patras
- 2005** MSc in Applications in Medical Sciences, Medical School, University of Patras

POSITIONS HELD

- Since 2015** Group Leader, Institute of Molecular Biology (IMB), Mainz
- 2013 – 2014** NIH Research Fellow, National Cancer Institute, National Institutes of Health (NIH), Bethesda
- 2008 – 2013** Postdoc, National Cancer Institute, National Institutes of Health (NIH), Bethesda

GROUP MEMBERS

- Marta Cipinska** PhD Student; since 05/2015
- Henrike Gothe** PhD Student; since 05/2016
- Monika Kuban** Lab Manager; since 01/2017
- Vera Minneker** PhD Student; since 04/2016
- Rossana Piccinno** PhD Student; since 04/2015
- Gabrielle Sant** PhD Student; since 11/2018

OVERVIEW

Maintaining the integrity of genetic information is essential for cell survival. Mechanisms that counteract DNA damage are important to help maintain cellular homeostasis by suppressing mutagenic events and genome rearrangements that may lead to disease, particularly cancer. One of the most severe forms of genome rearrangements are chromosome translocations. Translocations form by the illegitimate joining of chromosome breaks and often play key roles in the initial steps of tumorigenesis. Despite their prevalence and importance, our understanding of their genesis is, however, still rudimentary. Which are the molecular features that define recurrent chromosome breakpoints? How do the broken chromosome ends find each other within the nuclear space? Which are the DNA repair mechanisms that mediate the chromosome fusion and which are the factors that favour the interchromosomal fusion (translocation) over the intrachromosomal repair? By using a combination of molecular biology techniques, genetics and high-throughput imaging and sequencing approaches, we aim to shed light on the basic molecular mechanisms underlying the formation of oncogenic chromosome translocations.

RESEARCH HIGHLIGHTS

Novel imaging-based tools to probe rare, cancer-initiating genome rearrangements

Modelling the formation of recurrent cancer-initiating genome rearrangements of interest requires a versatile approach that can probe rare events with high sensitivity. We have now established a methodology, C-Fusion 3D, that uses fluorescence *in situ* hybridisation (FISH) to probe the position of individual chromosome ends of potential translocation partners in interphase cells in 3D. High-throughput microscopy and automated image analysis is

then used to identify single cells with chromosome breakage and translocations. This methodology complements existing approaches and offers several advantages in detection and quantification of translocations. It is: (a) suitable for detection of translocations without the requirement to map the precise translocation breakpoints or fusion product; (b) compatible with both site specific induction of breaks (mediated by endonucleases, ZNFs, CRISPR) and more physiological methods of inducing DNA damage, such as ionising radiation and chemotherapeutics (see below); (c) efficient in detecting translocations in interphase cells without the need of metaphase spread preparation with frequencies down to 10^{-4} . C-Fusion 3D is a powerful tool that can be used to dissect molecular and cellular mechanisms that contribute to the formation of any oncogenic chromosome translocation of interest.

Mechanistic insights into the formation of therapy-related, oncogenic translocations

A major problem following successful chemotherapy is the appearance of a second primary cancer, which is a direct consequence of the treatment due to the formation of cancer-initiating




Figure 1. Treatment with chemotherapeutics, such as alkylating agents or topoisomerase poisons has been associated with increased risk of developing secondary, therapy-induced cancers, driven by recurrent chromosome translocations.

translocations. As an example, therapy-induced acute myeloid leukaemias (t-AMLs) often develop after treatment with alkylating agents or topoisomerase-inhibitors and are characterised by distinct chromosome abnormalities that drive the occurrence of the secondary malignancies (Figure 1). Sequencing analysis of translocation junctions found in patients revealed recurrent fusion hotspots, indicative of defined molecular mechanisms that contribute to their formation. To identify molecular features that define recurrent breakpoint sites upon treatment with chemotherapeutic agents, we are using state-of-the-art genomic methodologies that map double-strand breaks (DSBs) across the genome with single nucleotide resolution. We then compare the high-resolution break-enrichment maps of chemotherapeutics with genomic, chromatin and topological features and associate recurrent break patterns with translocation hotspots. In combination with C-Fusion 3D, our efforts are focusing on: (1) identifying cellular and molecular pathways that contribute to the formation of recurrent DNA breaks at translocation hotspots found in patients; (2) understanding how the chromatin environment may predispose susceptibility to breakage and chromosome translocations; (3) assessing the influence of 3D genome organisation on chromosome breakage and fusion formation; and (4) identifying molecular players of the DNA damage response and novel factors that promote or inhibit the illegitimate fusion of chromosomes.

FUTURE DIRECTIONS

We will extend our analysis to identifying mechanisms governing the formation of different chromosome translocations leading to a variety of secondary malignancies. Central to our focus is shedding light on the events leading to the genesis of recurrent chromosome translocations involving the mixed lineage leukaemia (*MLL*) gene, which frequently occur upon treatment with the topoisomerase poison Etoposide. Moreover, in an effort to link DNA fragility with cellular physiology, we plan to profile endogenous DSBs across the genome in different cell types, with the

aim to identify common or cell type-specific signatures of DNA fragility. We will also develop novel tools that allow us to perform siRNA and CRISPR-based, unbiased and targeted screens to identify novel factors that govern key steps of the formation of translocations. Taken together, our research will shed light on the mechanisms of cancer-initiating translocations, which will advance our knowledge of the fundamental principles in cancer aetiology.

SELECTED PUBLICATIONS

Roukos V. (2018). Actin proteins assemble to protect the genome. *Nature*, 559: 35–37.

Roukos V and Misteli T. (2014). The biogenesis of chromosome translocations. *Nat Cell Biol*, 16: 293–300.

Roukos V, Voss TC, Schmidt CK, Lee S, Wangsa D and Misteli T. (2013). Spatial dynamics of chromosome translocations in living cells. *Science*, 341: 660–664.

NATALIA SOSHNIKOVA

“WE INVESTIGATE HOW
STEM CELLS ARE BORN”

EDUCATION

- 2004** PhD in Molecular Biology, Humboldt University, Berlin
1998 Diploma in Molecular Biology, Novosibirsk State University

POSITIONS HELD

- Since 2012** Group Leader, Institute of Molecular Biology (IMB), Mainz
2004 – 2011 Postdoc, University of Geneva

GROUP MEMBERS

- Jakub Klassek** Technician; 08/2016 – 08/2018
Michael Korzhev Staff Scientist; 03/2018 – 12/2018
Vadym Rostovskii Master's Student; 03/2018 – 12/2018
Nicholas Schön Master's Student; 03/2018 – 12/2018

OVERVIEW

In the adult organism, millions of cells are replaced every day. Adult stem cells are essential for the maintenance and regeneration of our complex organs. Yet, with age or in pathological conditions adult stem cells cease functioning properly. Understanding the natural path of adult stem cell specification during embryogenesis would allow for the generation of patient-specific adult stem cells. These could either be used for transplantation or for the screening of pharmacological compounds to develop personalised treatments. We use the mouse small intestine as a model system to decipher mechanisms of the intestinal stem cell (ISC) specification. We aim to identify signals and cells that trigger specification of embryonic ISCs. Moreover, we determine the degree of heterogeneity within the embryonic epithelial progenitors. Does their molecular heterogeneity reflect an early determination of distinct stem cell sub-types with potentially distinct functions? To address our questions, we employ mouse genetics, single-cell genomics and *ex vivo* 3D organoid assays.

RESEARCH HIGHLIGHTS

Until recently, the identity of embryonic epithelial cells giving rise to ISCs was unknown. In contrast to the morphologically distinct cells of the adult intestine, the embryonic small intestine is composed of morphologically identical, highly proliferative epithelial cells. Furthermore, adult ISCs can be identified based on the expression of a specific set of genes, so-called ISC signature genes. However, cells within the embryonic intestinal epithelium do not express these genes until 13.5 days of embryonic development (E13.5).

Using genetic cell fate mapping analysis we found that adult ISCs originate from molecularly distinct embryonic progenitors. For example, duodenal ISCs are specified from embryonic progenitors expressing *Foxa2*, whereas ISCs of the jejunum and

ileum come from cells expressing *Axin2*. Moreover, these distinct embryonic progenitors have different capacities to generate adult ISCs. This suggests that ISC progenitors are molecularly heterogeneous and might be different not only on the transcriptional but also on the functional level. To test this hypothesis, we performed single-cell RNA-sequencing analysis of the small intestines from mouse embryos. Based on the expression of markers and our genetic lineage tracing analysis, we separated the embryonic ISC progenitors into several groups. Currently, we are establishing a functional role for this heterogeneity during epithelial homeostasis and cancer.

To date, the signalling molecules triggering specification of the embryonic Lgr5⁺ ISC progenitors remain unknown. Wnt/ β -catenin signalling is a crucial regulator of adult ISCs maintenance and proliferation. Ablation of *β -catenin* or the Wnt receptors,

Lrp5/6, in the embryonic intestinal epithelium leads to reduced cell proliferation and a loss of ISC progenitors at E15.5. Further genetic experiments demonstrated that the secretion of Wnt ligands from mesenchymal cells is required for the maintenance of embryonic ISC progenitors at E15.5. Before this developmental stage, the expression of *Wnt* genes (that activate β -catenin dependent signalling) was not detected in either the embryonic intestinal epithelium or mesenchyme.

BMP signalling negatively controls self-renewal of adult ISCs and attenuates intestinal tumour formation. During embryonic development, several *Bmp* genes are expressed in either the intestinal mesenchyme or epithelium. We found that the *Id2* transcription factor, which is a target of TGF- β /BMP signalling, controls the timing of Lgr5⁺ progenitor specification during embryogenesis. In *Id2*-deficient embryos, Lgr5⁺ cells appear several days earlier and are much more abundant. Ectopic expression of Wnt ligands is required for the precocious activation of Wnt/ β -catenin target genes and earlier specification of Lgr5⁺ ISC progenitors in *Id2* mutant intestine. To identify factors essential for the specification of the embryonic ISC progenitors we analysed our single-cell RNA-sequencing data for the expression of various signalling molecules, including *Rspo*, Wnt, *Bmp/Tgf- β* and *Jag*, along with their receptors (Figure 1). Using mouse genetics and organoid assays we validated a function for these pathways during specification of the embryonic ISC progenitors.




Figure 1. Visualising Rspo/Wnt signalling during mouse development. The expression pattern of *Rspo3* (blue) and its receptor *Lgr5* (pink) in a paraffin section of a mouse embryo at E13.5 Scale bar: 30 μ m.

FUTURE DIRECTIONS

Our immediate goal is to understand the relationship of intestinal stem cells to cancer. We will examine the mechanisms underlying intestinal epithelial progenitor heterogeneity and whether some of these heterogeneous cells are more likely to

become cancer forming later in life. Furthermore, our long-term plan is to understand how the gut as an organ is maintained as a whole. Specifically, how both neurons and immune cells inside of the intestine maintain and communicate with the epithelium.

SELECTED PUBLICATIONS

Dzama MM, Nigmatullina L, Sayols S, Kreim N and Soshnikova N (2017). Distinct populations of embryonic epithelial progenitors generate Lgr5⁺ intestinal stem cells. *Dev Biol*, 432: 258–264.

Kazakevych J, Sayols S, Messner B, Krienke C and Soshnikova N. (2017). Dynamic changes in chromatin states during specification and differentiation of adult intestinal stem cells. *Nucleic Acids Res*, 45: 5770–5784.

Nigmatullina L, Norkin M, Dzama MM, Messner B, Sayols S and Soshnikova N. (2017). *Id2* controls specification of Lgr5(+) intestinal stem cell progenitors during gut development. *EMBO J*, 36: 869–885.

HELLE ULRICH

“TAILOR-MADE UBIQUITIN
LIGASES GIVE US INSIGHT
INTO SIGNALLING VIA
UBIQUITIN LINKAGE”



EDUCATION

- 2004** Habilitation in Genetics, Philipps University Marburg
1996 PhD in Chemistry, University of California, Berkeley
1992 Diploma in Biology, Georg August University Göttingen

POSITIONS HELD

- Since 2018** Executive Director, Institute of Molecular Biology (IMB), Mainz
Since 2013 Scientific Director, Institute of Molecular Biology (IMB), Mainz
 Professor, Faculty of Biology, Johannes Gutenberg University Mainz
2004 – 2012 Group Leader, Clare Hall Laboratories, Cancer Research UK London Research Institute
2000 – 2004 Group Leader, Max Planck Institute for Terrestrial Microbiology, Marburg
1998 – 2000 Postdoc, Max Planck Institute for Biochemistry, Martinsried
1997 – 1998 Postdoc, Centre for Molecular Biology (ZMBH), Heidelberg

GROUP MEMBERS

- Evrydiki Asimaki** PhD Student; since 09/2018
Liliana Batista Postdoc; 02/2013 – 08/2018
Sonja Braun Research Assistant; since 02/2016
Karolin Eifler-Olivi Postdoc; 10/2015 – 06/2018
Néstor García-Rodríguez Postdoc; 05/2013 – 08/2018
Srijana Maharjan Master's Student; since 10/2018
Julia Müller Research Assistant; 10/2017
Stephanie Nick PhD Student; since 11/2014
Regina Otto Personal Assistant; since 09/2015
Vanessa Patronelli Student Assistant; 04/2017
Kirill Petriukov PhD Student; since 08/2018
Christian Renz Postdoc; since 08/2014
Axel Schäfer PhD Student; since 10/2017
Jie Shi PhD Student; since 10/2018
Annie Sriramachandran Postdoc; since 03/2015
Tina Strauch Master's Student; since 10/2018
Laura Tomini Lab Manager; since 01/2015
Maria Villagomez-Torres PhD Student; since 01/2018
Vera Tröster PhD Student; since 01/2017
Sabrina Wegmann PhD Student; 11/2013 – 10/2018
Hans-Peter Wollscheid Postdoc; since 09/2014
Ronald Wong Postdoc; since 04/2013
George Yakoub PhD Student; since 01/2015
Nicola Zilio Postdoc; since 08/2014

OVERVIEW

Ubiquitin, as a posttranslational protein modifier, affects most aspects of cellular metabolism by modulating the stability, localisation or molecular interactions of its targets. Its versatility as a signalling molecule derives in part from its ability to form polymeric chains of varying topologies. Ubiquitin receptors with a basal affinity for the substrate along with one or several ubiquitin-binding domains that recognise the modification in a more or less linkage-selective manner, usually mediate the biological effects of polyubiquitylation. Structural biology has provided detailed insight into the principles of ubiquitin recognition by such receptors, while a wide range of analytical tools, including mass spectrometry, linkage-selective affinity probes, and selective inhibitors have facilitated the characterisation of ubiquitin conjugates. However, in many cases, the relevance of a particular chain linkage for its biological function is still unclear. Our poor understanding arises largely from a lack of suitable tools to directly manipulate the ubiquitin chain linkage. We have approached this problem by a new strategy of *in vivo* polyubiquitin chain engineering that involves the design and application of tailor-made ubiquitin protein ligases (E3s) for the purpose of “linkage mutagenesis”.

RESEARCH HIGHLIGHTS

Polyubiquitin chain linkage is controlled by the combination of the ubiquitin-conjugating enzymes (E2s) and E3s contributing to their assembly and is, therefore, an invariable feature of any ubiquitylation event. The experimental system underlying our approach is based on a well-defined K63-linked chain assembled on the replication factor PCNA in response to replication stress in budding yeast. Modification of PCNA at a single invariant lysine facilitates DNA replication in the presence of lesions and thereby contributes to genome stability. Monoubiquitylation by a specific E2-E3 pair promotes a mutagenic pathway of damage

bypass called translesion synthesis (TLS). Alternatively, extension of the modification to a K63-polyubiquitin chain by a dedicated E3, Rad5, initiates an alternative, error-free pathway known as template switching (TS). While the mechanism of TLS activation by PCNA monoubiquitylation is well understood, the function of the polyubiquitin chain in the TS pathway is still a matter of debate.

In order to examine the relevance of the K63-linkage in the context of DNA damage bypass, we designed three tailor-made E3s with defined, but distinct linkage specificities to replace Rad5 (Figure 1A). Selectivity for PCNA as a substrate was achieved by means of a conserved PCNA-interacting peptide (PIP), and well-characterised domains from unrelated E3s were used to install K63-, K48- and M1-specificity. Using a fusion of monoubiquitin to PCNA as a surrogate substrate, we found that in combination with the appropriate E2, all PIP-E3s generated PCNA-linked polyubiquitin chains *in vitro* (Figure 1B). The intended specificities were confirmed by linkage-specific antibodies or by using relevant ubiquitin mutants. Moreover, we found that the enzymes were E2- and substrate-specific and required prior monoubiquitylation of PCNA for activity. Kinetic analysis revealed substrate-binding properties very similar to Rad5 itself but with a range of different catalytic rate constants (Figure 1B). Expression of the PIP-E3s in a yeast *rad5Δ* mutant (Figure 1C)

and monitoring of DNA damage sensitivity was used to assess their *in vivo* function in the TS pathway (Figure 1D). Whereas the K63-specific E3 fully complemented the damage sensitivity of a *rad5Δ* mutant, PIP-E3(48) conferred a dominant negative effect that was attributable to an inhibition of the TLS pathway. We were able to show that this was due to proteasomal degradation of monoubiquitylated PCNA after its K48-linked polyubiquitylation. Surprisingly, the M1-specific E3 partially restored damage resistance to *rad5Δ*, and this was indeed mediated by PCNA modification and activation of TS. The partial nature of this effect is consistent with the lower intrinsic activity of this E3. Thus, our study has revealed that polyubiquitin chains of distinct linkages can sometimes substitute for each other, but can also convey alternative fates to the modified protein. At the same time, it is the first example of an *in vivo* manipulation of polyubiquitin chain linkage that illustrates a novel approach to analysing ubiquitin signalling in a physiological context.




Figure 1. Tailor-made E3s provide insight into the function of polyubiquitin chain linkage in DNA damage bypass. A) Linkage-specific polyubiquitylation of monoubiquitylated PCNA (Ub-PCNA) by designed PIP-E3s *in vitro*, detected by anti-PCNA immunoblotting. B) Kinetic parameters of PCNA diubiquitylation reactions, determined by fitting of initial rates to Michaelis Menten kinetics. C) Expression of PIP-E3(1) in *S. cerevisiae* (Hta2: histone H2A). D) Distinct effects of PIP-E3s on the sensitivity of *rad5Δ* cells towards an alkylating agent, methyl methanesulphonate (MMS).

FUTURE DIRECTIONS

Our strategy of linkage manipulation by tailor-made E3s would lend itself to the development of a promising and generally applicable research tool if the problems of substrate recognition and chain initiation can be overcome. While substrate recognition could, in principle, be generalised by replacing the PIP-PCNA interaction with an inducible dimerisation system, efficient chain initiation is more challenging, as our E3s were designed to only extend existing ubiquitin moieties on the substrate. In the case of PCNA, the proximal ubiquitin moiety is attached by a distinct

E2-E3 pair, which bypasses the need for direct modification of the substrate. In order to generalise this feature, we are currently developing a universal ubiquitin acceptor site that can be fused to the substrate of choice in combination with a dimerisation domain that would provide an interaction site for the E3. In this manner, we hope to generate a molecular tool analogous to existing degron systems that would allow the induced, linkage-specific polyubiquitylation of appropriately tagged substrates *in vivo*.

SELECTED PUBLICATIONS

García-Rodríguez N, Morawska M, Wong RP, Daigaku Y and Ulrich HD. (2018). Spatial separation between replisome- and template-induced replication stress signaling. *EMBO J*, 37: e98369.

García-Rodríguez N, Wong RP and Ulrich HD. (2018). The helicase Pif1 functions in the template switching pathway of DNA damage bypass. *Nucleic Acids Res*, 46: 8347–8356.

Hung SH, Wong RP, Ulrich HD[#] and Kao CF[#]. (2017). Monoubiquitylation of histone H2B contributes to the bypass of DNA damage during and after DNA replication. *Proc Natl Acad Sci USA*, 114: E2205–E22

[#] indicates joint correspondence

EVA WOLF

“WE STUDY PROTEIN INTERACTIONS INVOLVED IN CIRCADIAN REGULATION”



EDUCATION

- 2007** Habilitation in Biochemistry, Ruhr University, Bochum
1996 PhD in Biology, European Molecular Biology Laboratory (EMBL), Heidelberg
1991 Diploma in Biology, University of Heidelberg

POSITIONS HELD

- Since 2013** Adjunct Director, Institute of Molecular Biology (IMB), Mainz
 Professor of Structural Chronobiology, Faculty of Biology, Johannes Gutenberg University Mainz
2012 – 2013 Group Leader, Ludwig Maximilian University (LMU), Munich
2009 – 2011 Group Leader, Max Planck Institute for Biochemistry, Martinsried
2000 – 2009 Group Leader, Max Planck Institute for Molecular Physiology, Dortmund
1996 – 2000 Postdoc, Rockefeller University, New York

GROUP MEMBERS

- Arne Börgel** PhD Student; since 01/2016
Archit Garg PhD Student; since 11/2014
Tim Grimmelsmann PhD Student; since 01/2015
Silke Helmke Technician; since 05/2014
Shruti Krishnan PhD Student; since 10/2016
Torsten Merbitz-Zahradnik Postdoc; since 01/2014
Roberto Orru Postdoc; since 10/2015
Ruth Schaupp Personal Assistant, since 05/2014

OVERVIEW

Circadian clocks – operated by cell-autonomous transcription/translation feedback loops – affect many essential cellular, physiological and behavioural processes. In mammals, the transcription factors BMAL1/CLOCK activate three period (*per1,2,3*) and two cryptochrome (*cry1,2*) genes. The CRY1/2 and PER1/2 clock proteins repress BMAL1/CLOCK, whereas recruitment of co-activators to BMAL1/CLOCK leads to transcriptional activation. The mammalian Timeless (TIM) protein interacts with CRY but its role in the circadian clock is unclear. However, TIM together with its binding partner Tipin are implicated in DNA replication, checkpoint signalling and DNA repair. Hence, mammalian TIM is a multifunctional protein that may connect the circadian clock with genome maintenance by interacting with either CRY (circadian clock) or with replication fork associated proteins. To elucidate direct molecular links between circadian clocks and genome maintenance, we pursue 3D-structural, biochemical and quantitative biophysical analyses of complexes formed between CRY, TIM and Tipin and investigate their interplay within the circadian clock. These studies will advance our mechanistic understanding of how circadian gene regulation and genome maintenance are interconnected by time of day dependent changes in protein interaction networks.

RESEARCH HIGHLIGHTS

Accumulating evidence suggests a functional interconnection between mammalian circadian clocks and genome maintenance. For example, DNA damage induces phase advances of circadian rhythms, likely involving the Timeless (TIM) protein and ATM mediated damage signalling. Furthermore, Cryptochrome1 (CRY1) modulates ATR/Chk1-mediated DNA damage checkpoint responses in a circadian manner. To elucidate the molecular mechanisms underlying functional connections between the circadian clock and genome maintenance, we structurally, biochemically

and biophysically analyse interactions of the mammalian clock proteins CRY1/2 with TIM and the Timeless-interacting protein Tipin. Additionally, we analyse interactions of TIM and Tipin with replication fork associated proteins such as RPA.

We found, that the purified TIM-Tipin complex directly interacts with CRY1 and CRY2. The TIM-Tipin-CRY complex may recruit CRY proteins to the replication fork, for example, to modulate checkpoint responses. Conversely, the TIM-Tipin complex may be recruited to the circadian clock by its interaction with CRY1 or CRY2. While TIM, Tipin and the TIM-Tipin complex are known to be involved in checkpoint signalling and fork protection (Figure 1), Tipin and the TIM-Tipin complex have so far not been implicated in circadian regulation. Hence, the biological role of the mammalian TIM-Tipin-CRY complex in the circadian clock is unclear. We observed that CRY binding to the purified TIM-Tipin complex, to BMAL1 or PERIOD (PER) is mutually exclusive,

implying overlapping binding sites. We, therefore, speculate that the TIM-Tipin-CRY or TIM-CRY complex impacts on the circadian BMAL1/CLOCK transcriptional activity by interfering with the formation of PER-CRY or BMAL1-CRY complexes in the repressive phase of the circadian oscillator (Figure 1).

We further set out to identify molecular regions involved in TIM-Tipin-CRY interactions. We identified a short region in the mammalian TIM protein that modulates interactions with our purified CRY1 and CRY2 proteins. Deletion of this TIM region differentially affects CRY1 and CRY2 interactions *in vitro*, suggesting different biological roles of the TIM-(Tipin)-CRY1 and TIM-(Tipin)-CRY2 complexes. The identification of this TIM deletion construct now enables us to dissect the distinct roles of the TIM-(Tipin)-CRY1 and TIM-(Tipin)-CRY2 complexes in the circadian clock and in genome maintenance.

Using single particle Electron Microscopy (EM), we determined an initial negative stain EM structure from our purified TIM-Tipin-CRY complex samples. Compared to our previously reported TIM-Tipin-RPA Cryo-EM structure, the EM map of the TIM-Tipin-CRY sample has similar dimensions but lacks the regions that we assigned to RPA in the TIM-Tipin-RPA complex (Figure 1). We are currently pursuing 3D structural analyses of the TIM-Tipin-CRY complex at higher resolution. This will enable us to design additional mutants for the targeted functional analyses of TIM-Tipin-CRY and TIM-CRY interactions.




Figure 1. *Bottom:* Transcriptional regulation of the circadian clock in mouse liver. The transcription factor BMAL1/CLOCK is activated by coactivators during the day. BMAL1/CLOCK is repressed by a multi-subunit CRY/PER containing early repressor complex at night and later (morning) by CRY only. *Top:* TIM-Tipin (pink) in DNA replication and checkpoint signalling. Cryo-EM structure of the TIM-Tipin-RPA complex. *Middle:* negative stain EM envelope of a TIM-Tipin-CRY sample. The TIM-Tipin-CRY complex may connect the circadian clock and genome maintenance.

FUTURE DIRECTIONS

The roles of mammalian Timeless (TIM) in the circadian clock are vaguely defined and it is not known if or how the TIM-binding protein Tipin and the TIM-Tipin complex contribute to circadian regulation. Do TIM-CRY- or TIM-Tipin-CRY interactions affect circadian timing by altering CRY's activity as a transcriptional repressor of BMAL1/CLOCK? Furthermore, CRY and TIM are implicated in time of day dependent regulation of checkpoint signalling and advancing the circadian clock in phase in response to DNA damage. What are the underlying molecular mechanisms

and the distinct roles of TIM-CRY- or TIM-Tipin-CRY interactions in these processes? Finally, it will be interesting to find out if TIM-CRY- or TIM-Tipin-CRY interactions play a role in mammalian DNA repair pathways.

Our structural and biophysical protein-protein interaction analyses uncover molecule regions and amino acids involved in CRY-TIM-Tipin interactions. In future, this will enable us to design mutants and deletion constructs to address these interesting open questions in a targeted manner.


SELECTED PUBLICATIONS


Schmalen I, Reischl S, Wallach T, Klemz R, Grudziecki A, Prabu JR, Benda C, Kramer A and Wolf E. (2014). Interaction of circadian clock proteins CRY1 and PER2 is modulated by zinc binding and disulfide bond formation. *Cell*, 157: 1203–1215.

Witosz J, Wolf E and Mizuno N. (2014). Architecture and ssDNA interaction of the Timeless-Tipin-RPA COMPLEX. *Nucleic Acids Res*, 42: 12912–12927.

Czarna A, Berndt A, Singh HR, Grudziecki A, Ladurner AG, Timinszky G, Kramer A and Wolf E. (2013). Structures of *Drosophila* cryptochrome and Mouse cryptochrome1 provide insight into circadian function. *Cell*, 153: 1394–1405.

CORE FACILITIES





OVERVIEW

CORE FACILITIES

The Core Facilities at IMB provide access to state-of-the-art technology and offer services and training from experts.



There are currently seven Core Facilities (CFs) at IMB: Bioinformatics, Flow Cytometry, Genomics, Microscopy/Histology, Proteomics, Protein Production, and a Media Lab. The Bioinformatics, Genomics and Proteomics CFs provide users with a “full service”, from quality control of samples to data production and analysis. The Flow Cytometry and Microscopy/Histology CFs provide an “assisted service”, where researchers work independently on CF equipment after introductory training by CF staff. Whether receiving full or assisted service, the CFs’ staff are available for consultation and troubleshooting. Furthermore, CF staff often collaborate with researchers to provide customised or specialised services. For IMB researchers, all seven CFs are available for use. Beyond that, Flow Cytometry, Genomics, Microscopy/Histology and Proteomics CFs also offer their services to the larger research community in Mainz. CF services provided are based on user demand. For each facility, a user committee gives feedback on the equipment and user experience and helps define the services that each CF provides.

In addition to technical services, the CFs offer lectures on a variety of methods, as well as practical courses to instruct researchers in new techniques and instrumentation, data acquisition, experimental design, data processing, and analysis. These allow researchers to keep up-to-date with and broaden their knowledge of current and emerging technologies. Lectures are generally open to everyone.

Furthermore, IMB’s CFs are responsible for maintaining and providing training on core equipment that is available at IMB, as well as the radioactivity lab, the S2 lab, and IMB’s in-house animal facilities (mouse, zebrafish, *Xenopus* and *Drosophila*).

Andreas Vonderheit

Director of Core Facilities and Technology

BIOINFORMATICS

CORE FACILITY

The Bioinformatics Core Facility (BCF) supports researchers at IMB with computing infrastructure, web services, system administration, software training, and consulting on experimental design and statistics. In addition, BCF members actively participate in the computational processing, analysis, visualisation and interpretation of high-throughput "omics" data generated in the course of research projects.

SERVICES OFFERED

With their accumulated professional expertise, BCF staff offer support on different levels depending on project needs, ranging from basic IT and bioinformatics services to full-scale scientific collaborations in the context of "big data" research projects:

- + Consulting on the statistics and experimental design of genomics projects
- + Data quality assessment, processing, visualisation, interpretation and presentation of results
- + Development of analytical pipelines and their customisation for individual projects
- + Data mining of published datasets, correlation and integration of results
- + Assistance with the preparation of manuscripts, presentations and grant proposals
- + Workshops and tutorials on bioinformatics topics to facilitate data access and analysis
- + Testing, implementation and customisation of various software tools and online services
- + System administration and IT support in cooperation with the University of Mainz Data Center

The BCF operates a small computer cluster, a storage server, and web services such as Galaxy and R-Studio, which provide IMB researchers with a user-friendly interface to bioinformatics tools and databases. The Facility maintains a GitHub repository (github.com/imbforge) with software tools and pipelines dedicated to different types of next-generation sequencing (NGS) assays. BCF also offers customised solutions and long-term analytical support for numerous data-intensive IMB projects on a collaborative basis.



Emil Karaulanov Head

Since 10/2014

Anke Busch Bioinformatician

Since 01/2014

Nastasja Kreim Bioinformatician

Since 04/2012

Martin Oti Bioinformatician

Since 12/2017

Giuseppe Petrosino Bioinformatician

Since 03/2017

Christian Dietrich System Administrator

Since 04/2017

Pascal Silberhorn System Administrator

Since 12/2015




Figure 1. Time-course of RNA-Seq transcriptome profiling of human MCF10A cells. A) Principal component analysis plot of global sample similarity. B) Volcano plot depicting gene expression changes at a single time point with labelled top differential genes.

FLOW CYTOMETRY

CORE FACILITY

The Flow Cytometry Core Facility (FCCF) offers high-throughput measurements, analysis and separation of biological units through four different systems: a large particle sorter, a cell sorter, and two analysers. With this equipment, the FCCF can analyse and sort particles of 0.5 μm to 1,000 μm in diameter.

SERVICES OFFERED

The FCCF offers a full service for sorting and an assisted service along with training for the analysers. Additionally, its staff collaborates in terms of analysing flow cytometry data and sample preparation. During the past year, the FCCF has performed various types of experiments including multicolour measurements, cell separation for next generation sequencing, sorting of isolated neuronal nuclei, classical enrichments for subsequent cell culture, qPCR analysis, mass spectrometry, and microscopic investigations. Moreover, the FCCF performed reagent validation for the Protein Production Core Facility. The FCCF works with different types of material including: nuclei, stem cells, yeast, *C. elegans*, *Arabidopsis* seeds, and lipid droplets as well as various cultured cell lines and primary cells from humans, mice, zebrafish, and *Drosophila*. To educate and train users, the FCCF offers three different lectures per year, as well as an annual practical course for basic flow cytometry analysis and an advanced practical course for cell sorting.



Stefanie Möckel Head

Since 10/2016

Jesús Gil Pulido Staff Scientist

Since 09/2018

Nabil Boui Student Research Assistant

07/2018 – 11/2018

Ina Schäfer Biotechnologist

08/2011 – 06/2018




Figure 1. Analysis of apoptosis in Jurkat T cells treated with Camptothecin. Annexin V staining is used for analysis of exposed phosphatidylserine residues. Changes in mitochondrial membrane potential can be monitored with JC-1 staining. Both assays were performed in the FCCF's annual practical course.

GENOMICS

CORE FACILITY

The Genomics Core Facility (GCF) offers next-generation sequencing (NGS) services based on the Illumina NextSeq500 and MiSeq platforms. In 2018, the GCF acquired a MinIon sequencer (Oxford Nanopore).

SERVICES OFFERED

The GCF provides a full service for NGS, starting with the experimental design of the project and continuing up to the generation of sequencing data. In addition, the GCF also sequences self-prepared libraries from researchers at IMB.

After submission of RNA or DNA samples, the GCF performs initial quality control of the samples, library preparation, quality control of the prepared libraries, sequencing and raw data generation. Currently, the GCF supports library preparation for more than twenty applications as a standard service and develops new protocols to accommodate the user's needs for their specific projects. For example, in 2018, GCF supported members of Helle Ulrich's research group in establishing GLOE-Seq, a method for genome-wide detection of nicks, gaps, breaks and lesions.

The GCF also offers training in genomics techniques, including principles of NGS and the application of NGS in translational epigenetics.

RNA:

- + Strand specific mRNA-Seq, with poly-A selection
- + Strand specific total RNA-Seq, with rRNA depletion
- + Low input RNA-Seq
- + Small RNA-Seq
- + RIP-Seq
- + Bru-Seq
- + cDNA library preparation
- + circRNA
- + GRO-Seq
- + single-cell RNA (Smart-Seq2)
- + STARR-Seq

DNA:


- + ChIP-Seq
- + MBD-Seq
- + Whole genome sequencing
- + Whole genome bisulfite sequencing
- + Single-stranded DNA library preparation
- + Hi-C
- + MeDIP
- + 8-oxoG
- + DamID

User-prepared libraries:

- + iCLIP-Seq
- + Amplicon-Seq
- + ATAC-Seq
- + 4C / Capture-C
- + RR-MAB-Seq
- + GLOE-Seq
- + LAM-HGTS



Figure 1. NextSeq 500 benchtop, high throughput sequencer from Illumina.



Maria Mendez-Lago Head

Since 04/2016

Hanna Lukas Technician

Since 01/2013

Clara Werner Technician

Since 07/2015

MICROSCOPY & HISTOLOGY

CORE FACILITY

The Microscopy and Histology Core Facility (MHCF) provides state-of-the-art microscopes and histology instruments, as well as expertise in sample preparation and data post-processing. Users benefit from a broad range of lectures and hands-on training and can choose from independent, assisted, or full service.

SERVICES OFFERED

Microscopy users can select from 10 different instruments ranging from stereo microscopes and widefield microscopes to confocal, high-content screening and super-resolution microscopes. Four of the set-ups (one widefield, one scanning confocal, and two spinning disk confocal microscopes) are equipped for live cell imaging. Users are trained to work independently on the microscopes, although MHCF staff are always available to assist with sample preparation, image acquisition, as well as image processing, deconvolution and analysis (quantification). Beside licensed software programs for image deconvolution (Huygens Essential, SVI) and 3D visualisation/analysis (Imaris from Bitplane, Harmony from PerkinElmer), custom-made solutions are developed together with the users (e.g. by macro programming in open source software such as Fiji or ImageJ) or by the assembly of predefined building blocks in Columbus, a database and analysis software designed for high content imaging data (PerkinElmer). Super-resolution microscopy is offered as a full service or on a collaborative basis. User training for both microscopy and histology takes place throughout the year via practical courses and lectures, ranging in emphasis from confocal, live cell microscopy and super-resolution microscopy to image processing, as well as basics in histology and staining techniques.

In 2018, a new spinning disc microscope was co-funded by the DFG and put into operation. The system is devoted to live cell imaging of different samples. It is equipped with an ablation laser for the localised induction of DNA damage, a photomanipulation module for FRAP and TIRF, and two cameras for fast parallel imaging. Further, a 100TB server was installed to keep up with the increased demand for data storage and analysis on the Opera Phenix Screening Microscope.

For histology purposes, the MHCF provides a variety of histology techniques. In addition to semi-automated fixation and paraffin embedding, machines for the sectioning of paraffin-embedded tissue (microtome), frozen tissue (cryotome), and for gelatine/agarose embedded tissue and fresh tissue (vibratome) are part of the available instruments. Users may furthermore utilise optimised protocols for immunodetection, tissue clearing and solutions for classical tissue stainings (H&E, Masson Goldner Trichrome, PAS, and Azan).




Figure 1. 3D reconstruction of double smFISH against *dazl* (red) and *buc* (cyan) in a zebrafish stage II oocyte. They mark the germinal granules that will accumulate into germ plasma after fertilisation, a structure that is required for induction of primordial germ cell fate in the progeny and therefore fertility. Image acquired by Elke Roovers (Ketting group) with the spinning disk microscope "VisiScope" (Visitron, 40x water objective) at the IMB's Microscopy & Histology Core Facility.



Sandra Ritz Head

Since 01/2016

Mária Hanulová Staff Scientist

Since 02/2014

Jonas Schwirz Staff Scientist

Since 06/2016

PROTEIN PRODUCTION

CORE FACILITY

The Protein Production Core Facility (PPCF) provides support with the design, expression, purification and assay development of recombinant proteins that are in the focus of IMB's research. The facility also offers a variety of common protein tools that are routinely used by IMB researchers on a day-to-day basis.

SERVICES OFFERED

The PPCF supports researchers throughout the process of protein production. This includes the screening of suitable expression systems and vectors, optimisation of purification steps, upscaling of protein production and purification, as well as functional analysis and assay development with the purified products. The facility is equipped with two automated chromatography systems. These enable the use of the latest chromatographic methods required for state-of-the-art protein purification strategies.

Another key function of the PPCF is the generation and functional quality control of routine laboratory enzymes and affinity probes for IMB researchers. The facility is aiming to offer ready-to-use enzyme kits for all frequent applications at the institute.



Martin Möckel Head

Since 03/2018

Markus Matthes Head

02/2016 – 01/2018




Figure 1. Side view of the Äkta Prime Plus Chromatography system, showing inlet, injection and gradient valves, as well as the UV and conductivity monitor.

PROTEOMICS

CORE FACILITY

The Proteomics Core Facility (PCF) operates an EASY nLC 1000 ultraHPLC coupled online to a Q Exactive Plus mass spectrometer to perform proteomic measurements.

SERVICES OFFERED

As a general service, the PCF provides band identification, analysis of posttranslational modifications on single proteins and measurement of SILAC (stable isotope labelling with amino acids in cell culture) experiments. Additionally, the PCF also offers reductive dimethylation as a labelling technique and tandem mass tagging (TMT) for large-scale quantitation. TMT can be applied for quantifying up to 10 samples in parallel. In total 1,500 measurement hours annually are provided to IMB and the surrounding research centres in Mainz using a state-of-the-art mass spectrometry platform. The mass spectrometry service is provided as a full service, including initial consultation, sample preparation and basic proteomics data analysis by the PCF. Advanced proteomic workflows, label-free quantitation measurements, in-depth statistical and bioinformatics analysis are available in a collaborative context. The PCF offers lectures on proteomics and data analysis as well as providing researchers with hands-on experience during our practical courses.



Falk Butter Head

Since 05/2013

Jasmin Cartano Technician

Since 02/2014

Mario Dejung Bioinformatician

Since 05/2014

Anja Freiwald Engineer

Since 04/2013



Figure 1. The inside of an electrospray ionization (ESI) source showing ESI needle (upper right) pointing towards the mass spectrometry entrance. A fluid sample is transferred to the gaseous phase by the application of high voltage resulting in a fine spray. The same source housing can be modified for other ionization methods. Corona discharge needle (lower left) is necessary for atmospheric pressure chemical ionization (APCI).

MEDIA LAB

The Media Lab primarily supports scientific groups and other Core Facilities by producing media, buffers, and agar plates. In addition, the Media Lab is responsible for the administration of three supply centres, plasmid/cell line banks, general waste management and sterilisation of glassware.

SERVICES OFFERED

- + Supply of routinely-used buffers, solutions, liquid media and agar plates for molecular biological research and for culturing bacterial, yeast, and insect cells, as well as *C. elegans*
- + Production of made-to-order media
- + Management of three supply centres for enzymes, kits, and cell culture media
- + Administration of a vector data bank, human ORF clone collection and cell line bank
- + Overnight cultures for plasmid preparation
- + Sterilisation of solutions/media
- + Cleaning and sterilisation of glassware and lab equipment
- + Autoclaving of S1/S2 waste
- + Maintenance of in-house transport system




Figure 1. The media lab produces approximately 1,500 agar plates per week of different sizes and compositions depending on the needs of the user.



Andrea Haese-Corbit Head

Since 01/2018

Doris Beckhaus Assistant

Since 05/2011

Alwina Eirich Assistant

Since 07/2013

Pascal Hageböling Assistant

Since 01/2015

Annette Holstein Assistant

Since 04/2012

Marion Kay Assistant

Since 04/2016

Johann Suss Assistant

Since 04/2011



FACTS & FIGURES

IMB STAFF
PAGE 54



EXTRAMURAL
FUNDING
PAGE 55

RESEARCH
AND TRAINING
PAGE 56



TRAINING COURSES
PAGE 58

INVITED SPEAKERS
PAGE 60



AWARDS
PAGE 61

INTERNATIONAL
SCIENTIFIC EVENTS
PAGE 62



PUBLICATIONS
PAGE 63

SCIENTIFIC ADVISORY
BOARD
PAGE 71



RESEARCH ENVIRONMENT
PAGE 72

CAMPUS MAP
AND CONTACT
PAGE 74




IMPRINT
PAGE 77

IMB STAFF




IMB STAFF NATIONALITIES



EMPLOYEES BY STAFF CATEGORY

Group Leaders		6 %
Technical Staff		34 %
Administration & Scientific Management		11 %
PhD Students		33 %
Postdocs		16 %

EXTRAMURAL FUNDING



MAJOR FUNDERS



Boehringer Ingelheim Foundation (BIS)



German Research Foundation (DFG)



European Research Council (ERC)



Marie Curie Actions

FURTHER SUPPORT



Boehringer Ingelheim Fonds (BIF)



Federal Ministry of Education and Research (BMBWF)



German Academic Exchange Service (DAAD)



German Cancer Aid



European Molecular Biology Organisation (EMBO)



Joachim Herz Foundation



Natural Science Medical Research Center (NMFZ)



RESEARCH AND TRAINING

IMB is a thriving international research centre in Mainz. Our researchers explore science at the cutting edge of gene regulation, genome stability and other aspects of nuclear biology, ranging from the structural and molecular to the organismic level.

Currently, IMB supports 18 research groups with around 90 PhD students and 50 postdocs coming from 28 countries across the globe. This international environment has allowed ideas and expertise to be shared across different research priorities and has resulted in over 350 publications with 81 in 2018 alone. The individual research topics pursued at IMB centre on key questions in chromatin and RNA biology; epigenetics; DNA damage and genome stability; protein homeostasis; and related biomedical fields. These areas are producing exciting results that are transforming our understanding of how we develop and adapt to our environment, and how we age or develop disease.

IMB aims to ensure the best possible environment for its scientists by providing training in core competencies and professional skills. We thereby enable our researchers to tackle ambitious projects and develop their careers. A key part of the support offered comes through IMB's Core Facilities. Each facility is staffed by experts who advise, train and assist our researchers in the techniques and skills needed for cutting-edge science. Moreover, scientists at IMB are offered a wide range of professional skills courses that help push their competencies at all stages of their career. Courses offered cover topics such as presentation skills, scientific writing, project management, fundraising, career development, negotiation skills, and global leadership.



INTERNATIONAL PHD PROGRAMME

PhD students are a key part of our research at IMB. All of them participate in the International PhD Programme (IPP) on "Gene Regulation, Epigenetics & Genome Stability". This very successful programme was established with generous funding from the Boehringer Ingelheim Foundation and is supported by the participating institutions: IMB, Johannes Gutenberg University Mainz and the University Medical Center Mainz. Students benefit from the regular scientific exchange and networking across 46 research groups. In addition, they receive state-of-the-art scientific and professional skills training to support their personal development as a scientist.

www.imb.de/PhD

INTERNATIONAL POSTDOC PROGRAMME


IMB's Postdoc Programme (IPPro) has been established to meet the specific needs of postdocs, and to ensure that they are able to build the strongest possible foundation for success in their future careers. The programme provides sound scientific training through a variety of lectures, workshops and events, and offers postdocs full support with raising funds for their research. In addition to the guidance given by group leaders, the Postdoc Programme also offers mentoring discussions with IMB's scientific directors and invited speakers, career events, and preparation for applications and interviews. To succeed in today's competitive job market, postdocs must have excellent presentation, writing, project management and time management skills. As such, IMB provides courses and lectures on these topics. The programme also organises company site visits so that IMB postdocs have the opportunity to learn more about a range of future career opportunities.

www.imb.de/postdocs

INTERNATIONAL SUMMER SCHOOL

IMB's International Summer School (ISS) is a six-week programme for outstanding and enthusiastic undergraduate and Masters' students. Research groups participating in the ISS include group leaders at IMB, Johannes Gutenberg University and Mainz's University Medical Center. The ISS offers an attractive framework for training prospective scientists in an informal and international atmosphere. Lectures give students comprehensive insights into the latest research findings and identify key open questions in gene regulation, epigenetics and genome stability. Furthermore, the ISS offers courses on complementary skills, such as presentation and communication techniques, that are required for a successful career as a scientist. Most importantly, each student works on a cutting-edge research project within the lab of one of the participating research groups.

www.imb.de/ISS



TRAINING COURSES

In 2018, IMB offered the following training courses in scientific, technical and professional skills



LECTURES

CORE FACILITY	DATE	TITLE
GENERAL	14 May	Introduction to Molecular & Biochemistry Techniques
	02 Jul	Nuclear Magnetic Resonance (NMR)
BIOINFORMATICS	28 May	Databases in Bioinformatics
FLOW CYTOMETRY	09 Apr	Advanced Flow Cytometry: Principles of Cell Sorting
	25 Jun	Flow Cytometry
	22 Oct	Flow Cytometry: Introduction I
	29 Oct	Flow Cytometry: Introduction II
GENOMICS	11 Jun	Genomics (NGS)
MICROSCOPY & HISTOLOGY	23 Apr	Introduction to Microscopy
	30 Apr	Microscopy: F-Techniques & Super-Resolution
	07 May	Histology & Fluorescent Labeling
	18 Jun	Electron Microscopy
	18 Jun	Image Manipulation: The Slippery Slope to Misconduct
PROTEOMICS	16 Apr	Proteomics
PROTEIN PRODUCTION	04 Jun	Protein Production & Crystallography

PRACTICAL COURSES

CORE FACILITY	DATE	TITLE
BIOINFORMATICS	11 Apr	Bioinformatics: Introduction to R (Part I)
	18 Apr	Bioinformatics: Introduction to R (Part II)
	25 Apr	Bioinformatics: Introduction to R (Part III)
	02 May	Bioinformatics: Introduction to R (Part IV)
	16 May	Bioinformatics: ChIP-seq & RNA-seq Analysis with R (Part I)
	30 May	Bioinformatics: ChIP-seq & RNA-seq Analysis with R (Part II)
FLOW CYTOMETRY	27 Jun	Bioinformatics: ChIP-seq & RNA-seq Analysis with GALAXY
	19 – 24 Apr	Advanced Flow Cytometry: Principles of Cell Sorting
MICROSCOPY & HISTOLOGY	12 – 23 Nov	Flow Cytometry: Practical Course
	26 – 28 Feb	Histology Crash Course
PROTEOMICS	26 – 29 Mar	Microscopy: Image Processing & Analysis Course
	05 – 07 Mar	Proteomics: Practical Course



LECTURES

DATE	TITLE
16 Apr – 02 Jul	Lecture Series: Modern Techniques in Life Sciences
06 – 15 Aug	Advanced Lectures on Gene Regulation, Epigenetics & Genome Stability
03 Sep	Good Scientific Practice
15 Oct	Reaching your Readers Better in the Digital Age: From Cognitive Psychology to Practical Measures
16 Oct 2018 – 05 Feb 2019	Lecture Series: Introduction to Epigenetics
30 Oct	Intellectual Property Rights in Life Sciences – Basics, Strategies & Specialties, Pitfalls & Troubleshooting for Scientists

PRACTICAL COURSES

DATE	TITLE
09 – 10 Jan	Career Planning & Job Application Training
16 – 17 Jan	Grant Writing
25 – 26 Jan	How to Design Scientific Figures
26 Jan	Proposal Writing
31 Jan – 01 Feb	Scientific Writing
06 – 07 Feb	Project Management for Postdocs
13 Feb	Proposal Writing
21 – 22 Feb	Time Management
06 – 07 Mar	Project Management with Certification 1a
08 – 09 Mar	Biostatistics
13 – 14 Mar	Project Management with Certification 1b
13 – 14 Mar	Leadership Skills
20 – 21 Mar	Professional Communication
20 – 21 Mar	Project Management with Certification 2a+b
22 – 23 Mar	Convincing Scientific Presentations
27 – 28 Mar	Presenting Scientific Data
30 Mar	Intellectual Property Rights
30 – 31 Mar	Conflict Management
20 & 27 Apr	How to Create One's Job
13 – 14 Jun	Introduction to Biostatistics
25 Aug	Time & Self-Management
04 – 05 Sep	Writing for the Public
06 – 07 Sep	Convincing Scientific Presentations
18 – 19 Oct	Professional Communication
22 – 23 Nov	Time & Self-Management
28 – 29 Nov	Scientific Writing
11 Dec	Proposal Writing

INVITED SPEAKERS



DATE	EVENT	SPEAKER	INSTITUTION	TITLE
08 Feb	Seminar	ALFONSO VALENCIA	Barcelona Super-computing Center	Networks in proteins, diseases and (epi)genetics
19 Feb	TechTalk	KLAUS HERICK	ChromoTek, Martinsried	Smaller is better – ChromoTek’s nanobodies for immunoprecipitation, ChIP, MS, and imaging
22 Feb	Seminar	NIELS MAILAND	University of Copenhagen	Ubiquitin- and SUMO-mediated signalling in genome stability maintenance
26 Feb	TechTalk	JÖRG HILDMANN	BD Biosciences, Heidelberg	Overview of flow cytometry techniques for cell cycle, proliferation, live/dead, DNA staining and more – an inspiration for apoptosis and cell activation applications
22 Mar	Seminar	MATTHIAS SOLLER	University of Birmingham	Orchestrating mRNA processing in sex determination, neuronal development and function
19 Apr	Seminar	CATHERINE DARGEMONT	Paris Diderot University	The journey of mRNA from chromatin to the nuclear pore complex
23 Apr	Seminar	UTZ FISCHER	University of Würzburg	RNA biology and its role in human disease
03 May	Seminar	JAN KORBEL	EMBL Heidelberg	From genomic variation to molecular mechanism
17 May	Seminar	JOP KIND	Hubrecht Institute, Utrecht	Untangling spatial genome organisation: Detailed insights from single-cell DamID
18 May	Seminar	STELLA HURTLEY	<i>Science magazine</i> , Cambridge	Science publishing and publishing in <i>Science</i>
28 May	Seminar	RONALD HAY	University of Dundee	Harnessing the PML-SUMO-RNF4 axis for cancer therapy
07 Jun	Seminar	ANA POMBO	Max Delbrück Center for Molecular Medicine (MDC), Berlin	Genome architecture: Mechanisms of 3D chromatin folding
05 Jul	Seminar	JOHANNES GRAUMANN	Max Planck Institute for Heart and Lung Research, Bad Nauheim	Towards the whole sample spectrum: Extending proteomics from tissue culture to clinical samples
13 Sep	Seminar	ANN EHRENHOFER-MURRAY	Humboldt University, Berlin	Nutritional control of eukaryotic translation by queuosine and Dnmt2
25 Oct	Seminar	GERNOT LÄNGST	University of Regensburg	Nucleosomes stabilise RNA-DNA triple helices in human cells
08 Nov	Seminar	GAËLLE LEGUBE	Center for Integrative Biology, Toulouse	Chromatin and chromosome dynamics at DNA double-strand breaks
22 Nov	Seminar	WENDY BICKMORE	University of Edinburgh	Gene regulation in the 3D nucleus

AWARDS



CHRISTOF NIEHRS

Advanced Grant (ERC)

To pursue research into epigenetic gene regulation via R-loops

HELLE ULRICH

Proof of Concept Grant (ERC)

To develop a tool that allows adding specific ubiquitin modifications to proteins in cells

NATALIA SOSHNIKOVA

Heisenberg Fellowship (DFG)

To continue her research into stem cell specification during gut development

CHIARA PAOLANTONI

(Roignant Group)

PhD Fellowship from the Boehringer Ingelheim Fonds (BIF),

For research into the molecular mechanisms of m⁶A in *Drosophila* neurogenesis

WALTER BRONKHORST

(Ketting Group)

Postdoctoral Fellowship from the Peter und Traudl Engelhorn Stiftung

For research into the regulation of phase transition during zebrafish germ cell development and insights this may provide for neurodegenerative diseases

INTERNATIONAL SCIENTIFIC EVENTS

ORGANISED BY IMB IN 2018



15 March

SYMPOSIUM GENE REGULATION IN EVOLUTION: FROM MOLECULAR TO EXTENDED PHENOTYPES

Scientific organisers: Joachim Burger, Falk Butter, Susanne Foitzik, Thomas Hankeln, Martin Kaltenpoth and René Ketting

Gene Regulation in Evolution 15 MARCH 2018

INVITED SPEAKERS

- Irene Gallego Romero
- Your Gild
- Laurent Keller
- Ludovic Orlando
- John Parsch
- Amy Toth
- Patricia Winkler

SCIENTIFIC ORGANISERS

- Joachim Burger
- Falk Butter
- Susanne Foitzik
- Thomas Hankeln
- Martin Kaltenpoth
- René Ketting

VENUE: Institute of Molecular Biology (IMB), Ackermannweg 4, Mainz University Campus
www.imb.de/2018genevo, events@imb.de

EMBO Workshop

RNA and genome maintenance: Cooperation and conflict management

10 – 13 October 2018 | Mainz, Germany

ORGANIZER

SPEAKERS

REGISTRATION

CONTACT

meetings.embo.org/event/18-rna-genome

10-13 October

EMBO WORKSHOP RNA AND GENOME MAINTENANCE: COOPERATION AND CONFLICT MANAGEMENT

Scientific organisers: Andrés Aguilera, Petra Beli, Fabrizio d'Adda di Fagnaga, Brian Luke and Helle Ulrich

15-16 November

IMB WORKSHOP MOLECULAR MECHANISMS OF CIRCADIAN CLOCKS

Scientific organisers: Achim Kramer and Eva Wolf

IMB WORKSHOP

Molecular Mechanisms of Circadian Clocks

15-16 November 2018

INVITED SPEAKERS

- Ueli Schibler
- Charles Weitz
- Salvador Aznar Benitah
- Michael Brunner
- David Gattfield
- Carla Green
- Katja Lamia
- Carrie Partch
- Akhilesh Reddy
- Maria Robles
- Christoph Schiermann
- Hiroki Ueda

SCIENTIFIC ORGANISERS

- Achim Kramer
- Eva Wolf

Institute of Molecular Biology (IMB)
Ackermannweg 4, 55124 Mainz, Germany
www.imb.de/2018clock, events@imb.de



PUBLICATIONS

ANDRADE

Alanis-Lobato G, Mier P and Andrade-Navarro MA. (2018). The latent geometry of the human protein interaction network. *Bioinformatics*, 34: 2826–2834.

Brüne D, Andrade-Navarro MA and Mier P. (2018). Proteome-wide comparison between the amino acid composition of protein domains and linkers. *BMC Res Notes*, 11: 117.

Härtner F, Andrade-Navarro MA and Alanis-Lobato G. (2018). Geometric characterisation of disease modules. *Appl Netw Sci*, 3: 10.

Knuckles P, Lence T, Haussmann IU, Jacob D, Kreim N, Carl SH, Masiello I, Hares T, Villaseñor R, Hess D, Andrade-Navarro MA, Biggiogera M, Helm M, Soller M, Bühler M and Roignant JY. (2018). Zc3h13/Flacc is required for adenosine methylation by bridging the mRNA-binding factor Rbm15/Spentito to the m6A machinery component Wtap/Fl(2)d. *Genes Dev*, 32: 415–429.

Krefting J, Andrade-Navarro MA and Ibn-Salem J. (2018). Evolutionary stability of topologically associating domains is associated with conserved gene regulation. *BMC Biol*, 16: 87.

BAUMANN

Páez-Moscoso DJ, Pan L, Sigauke RF, Schroeder MR, Tang W and Baumann P. (2018). Pof8 is a La-related protein and a constitutive component of telomerase in fission yeast. *Nat Commun*, 9: 587.

Tseng CK, Wang HF, Schroeder MR, and Baumann P. (2018). The H/ACA complex disrupts triplex in hTR precursor to permit processing by RRP6 and PARN. *Nat Commun*, 9: 5430.

BELI

Borisova ME, Voigt A, Tollenaere MAX, Sahu SK, Juretschke T, Kreim N, Mailand N, Choudhary C, Bekker-Jensen S, Akutsu M, Wagner SA and Beli P. (2018). p38-MK2 signaling axis regulates RNA metabolism after UV-light-induced DNA damage. *Nat Commun*, 9: 1017.

Dev H, Chiang TWW, Lescale C, de Krijger I, Martin AG, Pilger D, Coates J, ... Beli P ... , Galanty Y and Jackson SP. (2018). Shieldin complex promotes DNA end-joining and counters homologous recombination in BRCA1-null cells. *Nat Cell Biol*, 20: 954–965.

Mier P and Andrade-Navarro MA. (2018). Glutamine codon usage and polyQ evolution in primates depend on the Q stretch length. *Genome Biol Evol*, 10: 816–825.

Schulz J, Mah N, Neuenschwander M, Kischka T, Ratei R, Schlag PM, Castañós-Vélez E, Fichtner I, Tunn PU, Denkert C, Klaas O, Berdel WE, von Kries JP, Makalowski W, Andrade-Navarro MA, Leutz A and Wethmar K. (2018). Loss-of-function uORF mutations in human malignancies. *Sci Rep*, 8: 2395.

Talyan S, Andrade-Navarro MA and Muro EM. (2018). Identification of transcribed protein coding remnants within lincRNAs. *Nucleic Acids Res*, 46: 8720–8729.

Taškova K, Fontaine JF, Mrowka R and Andrade-Navarro MA. (2018). Evaluation of *in vivo* and *in vitro* models of toxicity by comparison of toxicogenomics data with the literature. *Methods*, 132: 57–65.

Theobald J, Ghanem A, Wallisch P, Banaeiyan AA, Andrade-Navarro MA, Taškova K, Haltmeier M, Kurtz A, Becker H, Reuter S, Mrowka R, Cheng X and Wöflf S. (2018). Liver-kidney-on-chip to study toxicity of drug metabolites. *ACS Biomater Sci Eng*, 4: 78–89.



Heidelberger JB, Voigt A, Borisova ME, Petrosino G, Ruf S, Wagner SA and Beli P. (2018). Proteomic profiling of VCP substrates links VCP to K6-linked ubiquitylation and c-Myc function. *EMBO Rep*, 19: e44754.

PUBLICATIONS

BUTTER

Almeida MV, Dietz S, Redl S, Karaulanov E, Hildebrandt A, Renz C, Ulrich HD, König J, Butter F and Ketting RF. (2018). GTSF-1 is required for formation of a functional RNA-dependent RNA Polymerase complex in *Caenorhabditis elegans*. *EMBO J*, 37: e99325.

Baldi S, Jain DS, Harpprecht L, Zabel A, Scheibe M, Butter F, Straub T and Becker PB. (2018). Genome-wide rules of nucleosome phasing. *Mol Cell*, 72: 661–672.e4.

Dal Magro C, Keller P, Kotter A, Werner S, Duarte V, Marchand V, Ignarski M, Freiwald A, Müller R-U, Dieterich C, Motorin Y, Butter F, Atta M and Helm M. (2018). A vastly increased chemical variety of RNA modifications containing a thioacetal structure. *Angew Chemie Int Ed*, 57: 7893–7897.

Harris CJ*, Scheibe M*, Wongpalee SP, Liu W, Cornett EM, Vaughan RM, Li X, Chen W, Xue Y, Zhong Z, Yen L, Barshop WD, Rayatpisheh S, Gallego-Bartolome J, Groth M, Wohlschlegel JA, Du J, Rothbart SB, Butter F# and Jacobsen SE#. (2018). A DNA methylation reader complex that enhances gene transcription. *Science*, 362: 1182–1186.

Kiweler N, Brill B, Wirth M, Breuksch I, Laguna T, Dietrich C, Strand S, Schneider G, Groner B, Butter F, Heinzel T, Brenner W and Krämer OH. (2018). The histone deacetylases HDAC1 and HDAC2 are required for the growth and survival of renal carcinoma cells. *Arch Toxicol*, 92: 2227–2243.

Pons M, Reichardt CM, Hennig D, Nathan A, Kiweler N, Stocking C, Wichmann C, Christmann M, Butter F, Reichardt S, Schneider G, Heinzel T, Englert C, Hartkamp J, Krämer OH and Mahendrarajah N. (2018). Loss of Wilms tumor 1 protein is a marker for apoptosis in response to replicative stress in leukemic cells. *Arch Toxicol*, 92: 2119–2135.

Reis H, Schwesb M, Dietz S, Janzen CJ and Butter F. (2018). TelAP1 links telomere complexes with developmental expression site silencing in African trypanosomes. *Nucleic Acids Res*, 46: 2820–2833.

Becker K, Bluhm A*, Casas-Vila N*, Dinges N*, Dejung M, Sayols S, Kretz C, Roignant JY, Butter F# and Legewie S#. (2018). Quantifying post-transcriptional regulation in the development of *Drosophila melanogaster*. *Nat Commun*, 9: 4970.

Charbonnel C, Rymarenko O, Da Ines O, Benyahya F, White CI, Butter F and Amiard S. (2018). The linker histone GH1-HMGA1 is involved in telomere stability and DNA damage repair. *Plant Physiol*, 177: 311–327.

Roovers EF, Kaaij LJT, Redl S, Bronkhorst AW, Wiebrands K, de Jesus Domingues AM, Huang HY, Han CT, Riemer S, Dosch R, Salvenmoser W, Grün D, Butter F, van Oudenaarden A and Ketting RF. (2018).

Tdrd6a regulates the aggregation of Buc into functional subcellular compartments that drive germ cell specification. *Dev Cell*, 46: 285–301.e9.

Winzi M, Casas Vila N, Paszkowski-Rogacz M, Ding L, Noack S, Theis M, Butter F and Buchholz F. (2018). The long noncoding RNA *lncR492* inhibits neural differentiation of murine embryonic stem cells. *PLOS One*, 13: e0191682.

* indicates joint contribution # indicates joint correspondence





CREMER

Chen SY, Bestvater F, Schaufler W, Heintzmann R and Cremer C. (2018). Patterned illumination single molecule localization microscopy (piSMLM): User defined blinking regions of interest. *Opt. Express*, 26: 30009–30020.

Cremer C, Szczurek A, Strickfaden H, Birk U, Cremer M and Cremer T. (2018). Der Zellkern – eine Stadt in der Zelle, Teil 2. *Biol unserer Zeit*, 48: 45–53.

Cremer T, Cremer M and Cremer C. (2018). The 4D nucleome: Genome compartmentalization in an evolutionary context. *Biochem*, 83: 313–325.

Eryilmaz M, Schmitt E, Krufczik M, Theda F, Lee JH, Cremer C, Bestvater F, Schaufler W, Hausmann M and Hildenbrand G. (2018). Localization microscopy analyses of MRE11 clusters in 3D-conserved cell nuclei of different cell lines. *Cancer*, 10: 25.

Kordon MM, Szczurek A, Berniak K, Szelest O, Solarczyk K, Tworzydło M, Wachsmann-Hogiu S, Vaahtokari A, Cremer C, Pederson T and Dobrucki JW. (2018). PML-like subnuclear bodies, containing XRCC1, juxtaposed to DNA replication-based single-strand breaks. *FASEB J*, doi: 10.1096/fj.201801379R.

Schink A, Neumann J, Leifke AL, Ziegler K, Fröhlich-Nowoisky J, Cremer C, Thines E, Weber B, Pöschl U, Schuppan D and Lucas K. (2018). Screening of herbal extracts for TLR2- and TLR4-dependent anti-inflammatory effects. *PLOS ONE*, 13: e0203907.

Szczurek A, Birk U, Knecht H, Dobrucki J, Mai S and Cremer C. (2018). Super-resolution binding activated localization microscopy through reversible change of DNA conformation. *Nucleus*, 9: 182–189.

KETTING

Roovers EF, Kaaij LJT, Redl S, Bronkhorst AW, Wiebrands K, de Jesus Domingues AM, Huang HY, Han CT, Riemer S, Dosch R, Salvenmoser W, Grün D, Butter F, van Oudenaarden A and Ketting RF. (2018). Tdrd6a regulates the aggregation of Buc into functional subcellular compartments that drive germ cell specification. *Dev Cell*, 46: 285–301.e9.

Kaaij LJT, van der Weide RH, Ketting RF and de Wit E. (2018). Systemic loss and gain of chromatin architecture throughout zebrafish development. *Cell Rep*, 24: 1–10.e4.

Almeida MV, Dietz S, Redl S, Karaulanov E, Hildebrandt A, Renz C, Ulrich HD, König J, Butter F and Ketting RF. (2018). GTSF-1 is required for formation of a functional RNA-dependent RNA Polymerase complex in *Caenorhabditis elegans*. *EMBO J*, 37: e99325.

Bronkhorst AW and Ketting RF. (2018). Trimming it short: PNLDC1 is required for piRNA maturation during mouse spermatogenesis. *EMBO Rep*, 19: e45824.

KHMELINSKII

Meurer M, Duan Y, Sass E, Kats I, Herbst K, Buchmuller BC, Dederer V, Huber F, Kirmaier D, Štefl M, Van Laer K, Dick TP, Lemberg MK, Khmelinskii A[#], Levy ED[#] and Knop M[#]. (2018). Genome-wide C-SWAT library for high-throughput yeast genome tagging. *Nat Methods*, 15: 598–600.

Kowalski L, Bragoszewski P, Khmelinskii A, Glow E, Knop M and Chacinska A. (2018). Determinants of the cytosolic turnover of mitochondrial intermembrane space proteins. *BMC Biol*, 16: 66.

Kats I, Khmelinskii A, Kschonsak M, Huber F, Knieß RA, Bartosik A and Knop M. (2018). Mapping degradation signals and pathways in a eukaryotic N-terminome. *Mol Cell*, 70: 488–501.



[#] indicates joint correspondence

PUBLICATIONS

KÖNIG

Almeida MV, Dietz S, Redl S, Karaulanov E, Hildebrandt A, Renz C, Ulrich HD, König J, Butter F and Ketting RF. (2018). GTSF-1 is required for formation of a functional RNA-dependent RNA Polymerase complex in *Caenorhabditis elegans*. *EMBO J*, 37: e99325.

Braun S*, Enculescu M*, Setty ST*, Cortés-López M, de Almeida BP, Sutandy FXR, Schulz L, Busch A, Seiler M, Ebersberger S, Barbosa-Morais NL, Legewie S#, König J# and Zarnack K#. (2018). Decoding a cancer-relevant splicing decision in the RON proto-oncogene using high-throughput mutagenesis. *Nat Commun*, 9: 3315.

LEGEWIE

Becker K, Bluhm A*, Casas-Vila N*, Dinges N*, Dejung M, Sayols S, Kretz C, Roignant JY, Butter F# and Legewie S#. (2018). Quantifying post-transcriptional regulation in the development of *Drosophila melanogaster*. *Nat Commun*, 9: 4970.

Braun S*, Enculescu M*, Setty ST*, Cortés-López M, de Almeida BP, Sutandy FXR, Schulz L, Busch A, Seiler M, Ebersberger S, Barbosa-Morais NL, Legewie S#, König J# and Zarnack K#. (2018). Decoding a cancer-relevant splicing decision in the RON proto-oncogene using high-throughput mutagenesis. *Nat Commun*, 9: 3315.

Fritsch C*, Baumgärtner S*, Kuban M, Steinshorn D, Reid G# and Legewie S#. (2018). Estrogen-dependent control and cell-to-cell variability of transcriptional bursting. *Mol Syst Biol*, 14: e7678.

LEMKE

Das DK, Govindan R, Nikić-Spiegel I, Krammer F, Lemke EA and Munro JB. (2018). Direct visualization of the conformational dynamics of single influenza hemagglutinin trimers. *Cell*, 174: 926–937.e12.

Fuertes G, Banterle N, Ruff KM, Chowdhury A, Pappu R V, Svergun DI and Lemke EA. (2018). Comment on "Innovative scattering analysis shows that hydrophobic disordered proteins are expanded in water." *Science*, 361: eaau8230.

Hellenkamp B, Schmid S, Doroshenko O, Opanasyuk O, Kühnemuth R, Rezaei Adariani S, Ambrose B, ... Lemke EA ..., Craggs TD and Hugel T. (2018). Precision and accuracy of single-molecule FRET measurements – a multi-laboratory benchmark study. *Nat Methods*, 15: 669–676.

Olgeiser L, Haag C, Boerner S, Ule J, Busch A, Koepke J, König J, Feldbrügge M and Zarnack K. (2018). The key protein of endosomal mRNP transport Rrm4 binds translational landmark sites of cargo mRNAs. *EMBO Rep*, 20: e46588.

Sutandy FXR*, Ebersberger S*, Huang L*, Busch A, Bach M, Kang HS, Fallmann J, Maticzka D, Backofen R, Stadler PF, Zarnack K, Sattler M, Legewie S# and König J. (2018). *In vitro* iCLIP-based modeling uncovers how the splicing factor U2AF2 relies on regulation by cofactors. *Genome Res*, 28: 699–713.

Ghosh B*, Sarma U*, Sourjik V# and Legewie S#. (2018). Sharing of phosphatases promotes response plasticity in phosphorylation cascades. *Biophys J*, 114: 223–236.

Strasen J*, Sarma U*, Jentsch M*, Bohn S, Sheng C, Horbelt D, Knaus P, Legewie S# and Loewer A#. (2018). Cell-specific responses to the cytokine TGFβ are determined by variability in protein levels. *Mol Syst Biol*, 14: e7733.

Sutandy FXR*, Ebersberger S*, Huang L*, Busch A, Bach M, Kang HS, Fallmann J, Maticzka D, Backofen R, Stadler PF, Zarnack K, Sattler M, Legewie S# and König J. (2018). *In vitro* iCLIP-based modeling uncovers how the splicing factor U2AF2 relies on regulation by cofactors. *Genome Res*, 28: 699–713.

Koehler C and Lemke EA. (2018). MultiBacTAG-Genetic code expansion using the baculovirus expression system in Sf21 Cells. pages 297-312, In: Noncanonical amino acids: *Methods in Molecular Biology*, Vol. 1728. Lemke EA (ed). *Humana Press*, New York.

Kormos A, Koehler C, Fodor EA, Rutkai ZR, Martin ME, Mezö G, Lemke EA and Kele P. (2018). Bistetrazine-cyanines as double-clicking fluorogenic two-point binder or crosslinker probes. *Chem - A Eur J*, 24: 8841–8847.

Kozma E, Paci G, Girona HE, Lemke EA and Kele P. Fluorogenic tetrazine-siliconrhodamine probe for the labeling of noncanonical amino acid tagged proteins. pages 337-363, In: Noncanonical amino acids: *Methods in Molecular Biology*, Vol. 1728. Lemke EA (ed). *Humana Press*, New York.

Lemke EA (ed) (2018). Noncanonical amino acids: *Methods in Molecular biology*, Vol. 1728. *Humana Press*, New York. 411 pages.

* indicates joint contribution # indicates joint correspondence



Mikhaleva S and Lemke EA. (2018). Beyond the transport function of import receptors: What's all the FUS about? *Cell*, 173: 549–553.

Tan PS, Aramburu IV, Mercadante D, Tyagi S, Chowdhury A, Spitz D, Shammass SL, Gräter F and Lemke EA. (2018). Two differential binding mechanisms of FG-nucleoporins and nuclear transport receptors. *Cell Rep*, 22: 3660–3671.

LUKE

Luke B. (2018). Telomere regulation (guest editor). *Differentiation*, 102: 27–29.

Michelini F, Jalihal AP, Francia S, Meers C, Neeb ZT, Rossiello F, Gioia U, Aguado J, Jones-Weinert C, Luke B, Biamonti G, Nowacki M, Storici F, Carninci P, Walter NG and d'Adda di Fagagna F. (2018). From “cellular” RNA to “smart” RNA: Multiple roles of RNA in genome stability and beyond. *Chem Rev*, 118: 4365–4403.

Misino S, Bonetti D, Luke-Glaser S and Luke B. (2018). Increased TERRA levels and RNase H sensitivity are conserved hallmarks of post-senescent survivors in budding yeast. *Differentiation*, 100: 37–45.

NIEHRS

Busceti CL, Di Menna L, Bianchi F, Mastroiacovo F, Di Pietro P, Traficante A, Bozza G, Niehrs C, Battaglia G, Bruno V, Fornai F, Volpe M, Rubattu S and Nicoletti F. (2018). Dickkopf-3 causes neuroprotection by inducing vascular endothelial growth factor. *Front Cell Neurosci*, 12: 292.

Colditz J, Thiele S, Baschant U, Niehrs C, Bonewald LF, Hofbauer LC and Rauner M. (2018). Postnatal skeletal deletion of Dickkopf-1 increases bone formation and bone volume in male and female mice, despite increased sclerostin expression. *J Bone Miner Res*, 33: 1698–1707.

Dietrich K, Baumgart J, Eshkind L, Reuter L, Godtel-Armbrust U, Butt E, Mushev M, Marini F, More P, Grosser T, Niehrs C, Wojnowski L and Mathäs M. (2018). Health-relevant phenotypes in the offspring of mice given CAR activators prior to pregnancy. *Drug Metab Dispos*, 46: 1827–1835.

Karamariti E, Zhai C, Yu B, Qiao L, Wang Z, Potter CMF, Wong MM, Simpson RML, Zhang Z, Wang X, del Barco Barrantes I, Niehrs C, Kong D, Zhao Q, Zhang Y, Hu Y, Zhang C and Xu Q. (2018). DKK3 (Dickkopf 3) alters atherosclerotic plaque phenotype involving vascular progenitor and fibroblast differentiation into smooth muscle cells. *Arterioscler Thromb Vasc Biol*, 38: 425–437.

Tan PS and Lemke EA. (2018). Probing differential binding mechanisms of phenylalanine-glycine-rich nucleoporins by single-molecule FRET. *Methods in Enzymology*, 611:327–346.

Wawryszyn M, Sauter PF, Nieger M, Koos MRM, Koehler C, Luy B, Lemke EA and Bräse S. (2018). Synthesis of azido-glycans for chemical glycomodification of proteins. *European J Org Chem*, 2018: 4296–4305.



Martins LR, Bung RK, Koch S, Richter K, Schwarzmüller L, Terhardt D, Kurtulmus B, Niehrs C, Rouhi A, Lohmann I, Pereira G, Fröhling S, Glimm H and Scholl C. (2018). Stk33 is required for spermatid differentiation and male fertility in mice. *Dev Biol*, 433: 84–93.

Melnik S, Dvornikov D, Müller-Decker K, Depner S, Stanek P, Meister M, Warth A, Thomas M, Muley T, Risch A, Plass C, Klingmüller U, Niehrs C and Glinka A. (2018). Cancer cell specific inhibition of Wnt/β-catenin signaling by forced intracellular acidification. *Cell Discov*, 4: 37.

Schäfer A, Mekker B, Mallick M, Vastolo V, Karaulanov E, Sebastian D, von der Lippen C, Epe B, Downes DJ, Scholz C and Niehrs C. (2018). Impaired DNA demethylation of C/EBP sites causes premature aging. *Genes Dev*, 32: 742–762.

PUBLICATIONS

RICHLY

Wilhelm T and Richly H. (2018). Evolution does not care. *TheScienceBreaker*, 4.

Kaymak A, Sayols S, Papadopoulou T and Richly H. (2018). Role for the transcriptional activator ZRF1 in early metastatic events in breast cancer progression and endocrine resistance. *Oncotarget*, 9: 28666–28690.

Wilhelm T and Richly H. (2018). Autophagy during ageing – from Dr Jekyll to Mr Hyde. *FEBS J*, 285: 2367–2376.

Chitale S and Richly H. (2018). H4K20me2: Orchestrating the recruitment of DNA repair factors in nucleotide excision repair. *Nucleus*, 9: 212–215.

Chitale S and Richly H. (2018). DICER- and MMSET-catalyzed H4K20me2 recruits the nucleotide excision repair factor XPA to DNA damage sites. *J Cell Biol*, 217: 527–540.

Medina R and Richly H. (2018). The dire side of autophagy in aging: Lessons from *C. elegans*. *Commun Integr Biol*, 11: e1395120.

ROIGNANT

Angelova MT, Dimitrova DG, Dinges N, Lence T, Worpenberg L, Carré C# and Roignant JY#. (2018). The emerging field of epitranscriptomics in neurodevelopmental and neuronal disorders. *Front Bioeng Biotechnol*, 6: 46.

Akhtar J, Kreim N, Marini F, Mohana GK, Brune D, Binder H and Roignant JY. (2018). The exon junction complex regulates the release and phosphorylation of paused RNA polymerase II. *bioRxiv*, doi: 10.1101/271544.

Becker K, Bluhm A*, Casas-Vila N*, Dinges N*, Dejung M, Sayols S, Kretz C, Roignant JY, Butter F# and Legewie S#. (2018). Quantifying post-transcriptional regulation in the development of *Drosophila melanogaster*. *Nat Commun*, 9: 4970.

de Brouwer* APM, Jamra* R, Körtel* N, Soyris C, Polla DL, Safra M, Zisso A, ... Roignant JY#, van Bokhoven H# and Schwartz S#. (2018). Variants in PUS7 cause intellectual disability with speech delay, microcephaly, short stature, and aggressive behavior. *Am J Hum Gen*, 6: 1045–1052.

Jantsch MF, Quattrone A, O'Connell M, Helm M, Frye M, Macias-Gonzales M, Ohman M, ... Roignant JY ..., Ule J and Fray R. (2018). Positioning Europe for the EPITRANSCRIPTOMICS challenge. *RNA Biol*, 15: 829–831.

Knuckles P, Lence T, Haussmann IU, Jacob D, Kreim N, Carl SH, Masiello I, Hares T, Villaseñor R, Hess D, Andrade-Navarro MA, Biggiogera M, Helm M, Soller M, Bühler M and Roignant JY. (2018). Zc3h13/Flacc is required for adenosine methylation by bridging the mRNA-binding factor Rbm15/Spentito to the m6A machinery component Wtap/Fl(2)d. *Genes Dev*, 32: 415–429.

Lence T, Paolantoni C, Worpenberg L and Roignant JY. (2018). Mechanistic insights into m⁶A RNA enzymes. *Biochim Biophys Acta Gene Regul Mech*, doi: 10.1016/j.bbagrm.2018.10.014.

ROUKOS

Gothe HJ, Minneker V and Roukos V (2018). Dynamics of double-strand breaks: Implications for the formation of chromosome Translocations. pages 27-38, In: Chromosome translocation: *Advances in Experimental Medicine and Biology*, Vol. 1044. Zhang Y (ed). *Springer*, Singapore.

Roukos V. (2018). Actin proteins assemble to protect the genome. *Nature*, 559: 35–37.

Zirkel A, Nikolic M, Sofiadis K, Mallm JP, Brackley CA, Gothe H, Drechsel O, ... Roukos V..., Rippe K and Papantonis A. (2018). HMGB2 loss upon senescence entry disrupts genomic organization and induces CTCF clustering across cell types. *Mol Cell*, 70: 730–744.e6.

Dörsam B, Seiwert N, Foersch S, Stroh S, Nagel G, Begaliew D, Diehl E, Kraus A, McKeague M, Minneker V, Roukos V, Reißig S, Waisman A, Moehler M, Stier A, Mangerich A, Dantzer F, Kaina B and Fahrer J. (2018). PARP-1 protects against colorectal tumor induction, but promotes inflammation-driven colorectal tumor progression. *Proc Natl Acad Sci USA*, 115: E4061–E4070.

* indicates joint contribution # indicates joint correspondence



SOSHNIKOVA

Szenker-Ravi E, Altunoglu U, Leushacke M, Bosso-Lefèvre C, Khatoo M, Thi Tran H, Naert T, ... Soshnikova N ..., Kayserili H and Reversade B. (2018). RSP02 inhibition of RNF43 and ZNRF3 governs limb development independently of LGR4/5/6. *Nature*, 557: 564–569.

ULRICH

Almeida MV, Dietz S, Redl S, Karaulanov E, Hildebrandt A, Renz C, Ulrich HD, König J, Butter F and Ketting RF. (2018). GTSF-1 is required for formation of a functional RNA-dependent RNA Polymerase complex in *Caenorhabditis elegans*. *EMBO J*, 37: e99325.

García-Rodríguez N, Morawska M, Wong RP, Daigaku Y and Ulrich HD. (2018). Spatial separation between replisome- and template-induced replication stress signaling. *EMBO J*, 37: e98369.

García-Rodríguez N, Wong RP and Ulrich HD. (2018). The helicase Pif1 functions in the template switching pathway of DNA damage bypass. *Nucleic Acids Res*, 46: 8347–8356.



CORE FACILITIES

Almeida MV, Dietz S, Redl S, Karaulanov E, Hildebrandt A, Renz C, Ulrich HD, König J, Butter F and Ketting RF. (2018). GTSF-1 is required for formation of a functional RNA-dependent RNA Polymerase complex in *Caenorhabditis elegans*. *EMBO J*, 37: e99325.

Becker K, Bluhm A*, Casas-Vila N*, Dinges N*, Dejung M, Sayols S, Kreutz C, Roignant JY, Butter F# and Legewie S#. (2018). Quantifying post-transcriptional regulation in the development of *Drosophila melanogaster*. *Nat Commun*, 9: 4970.

Borisova ME, Voigt A, Tollenaere MAX, Sahu SK, Juretschke T, Kreim N, Mailand N, Choudhary C, Bekker-Jensen S, Akutsu M, Wagner SA and Beli P. (2018). p38-MK2 signaling axis regulates RNA metabolism after UV-light-induced DNA damage. *Nat Commun*, 9: 1017.

Braun S*, Enculescu M*, Setty ST*, Cortés-López M, de Almeida BP, Sutandy FXR, Schulz L, Busch A, Seiler M, Ebersberger S, Barbosa-Morais NL, Legewie S#, König J# and Zarnack K#. (2018). Decoding a cancer-relevant splicing decision in the RON proto-oncogene using high-throughput mutagenesis. *Nat Commun*, 9: 3315.

Charbonnel C, Rymarenko O, Da Ines O, Benyahya F, White CI, Butter F and Amiard S. (2018). The linker histone GH1-HMGA1 is involved in telomere stability and DNA damage repair. *Plant Physiol*, 177: 311–327.

Dal Magro C, Keller P, Kotter A, Werner S, Duarte V, Marchand V, Ignarski M, Freiwald A, Müller R-U, Dieterich C, Motorin Y, Butter F, Atta M and Helm M. (2018). A vastly increased chemical variety of RNA modifications containing a thioacetal structure. *Angew Chemie Int Ed*, 57: 7893–7897.

Heidelberger JB, Voigt A, Borisova ME, Petrosino G, Ruf S, Wagner SA and Beli P. (2018). Proteomic profiling of VCP substrates links VCP to K6-linked ubiquitylation and c-Myc function. *EMBO Rep*, 19: e44754.

Kaymak A, Sayols S, Papadopoulou T and Richly H. (2018). Role for the transcriptional activator ZRF1 in early metastatic events in breast cancer progression and endocrine resistance. *Oncotarget*, 9: 28666–28690

Kiweler N, Brill B, Wirth M, Breuksch I, Laguna T, Dietrich C, Strand S, Schneider G, Groner B, Butter F, Heinzel T, Brenner W and Krämer OH. (2018). The histone deacetylases HDAC1 and HDAC2 are required for the growth and survival of renal carcinoma cells. *Arch Toxicol*, 92: 2227–2243.

Knuckles P, Lence T, Haussmann IU, Jacob D, Kreim N, Carl SH, Masiello I, Hares T, Villaseñor R, Hess D, Andrade-Navarro MA, Biggiogera M, Helm M, Soller M, Bühler M and Roignant JY. (2018). Zc3h13/Flacc is required for adenosine methylation by bridging the mRNA-binding factor Rbm15/Spentito to the m6A machinery component Wtap/Fl(2)d. *Genes Dev*, 32: 415–429.

PUBLICATIONS

Momper R, Steinbrecher J, Dorn M, Rörich I, Bretschneider S, Tonigold M, Ramanan C, Ritz S, Mailänder V, Landfester K and Bannwarth MB. (2018). Enhanced photoluminescence properties of a carbon dot system through surface interaction with polymeric nanoparticles. *J Colloid Interf Sc*, 518: 11–20.

Pons M, Reichardt CM, Hennig D, Nathan A, Kiweler N, Stocking C, Wichmann C, Christmann M, Butter F, Reichardt S, Schneider G, Heinzl T, Englert C, Hartkamp J, Krämer OH and Mahendrarajah N. (2018). Loss of Wilms tumor 1 protein is a marker for apoptosis in response to replicative stress in leukemic cells. *Arch Toxicol*, 92: 2119–2135.

Raden M, Ali SM, Alkhnabshi OS, Busch A, Costa F, Davis JA, Eggenhofer F, Gelhausen R, Georg J, Heyne S, Hiller M, Kundu K, Kleinkauf R, Lott SC, Mohamed MM, Mattheis A, Miladi M, Richter AS, Will S, Wolff J, Wright PR and Backofen R. (2018). Freiburg RNA tools: A central online resource for RNA-focused research and teaching. *Nucleic Acids Res*, 46: W25-W29.

Reinholz J, Diesler C, Schöttler S, Kokkinopoulou M, Ritz S, Landfester K and Mailänder V. (2018). Protein machineries defining pathways of nanocarrier exocytosis and transcytosis. *Acta Biomater*, 11: 432–443.

Reis H, Schwebs M, Dietz S, Janzen CJ and Butter F. (2018). TelAP1 links telomere complexes with developmental expression site silencing in African trypanosomes. *Nucleic Acids Res*, 46: 2820–2833.

Roovers EF, Kaaij LJT, Redl S, Bronkhorst AW, Wiebrands K, de Jesus Domingues AM, Huang HY, Han CT, Riemer S, Dosch R, Salvenmoser W, Grün D, Butter F, van Oudenaarden A and Ketting RF. (2018). Tdrd6a regulates the aggregation of Buc into functional subcellular compartments that drive germ cell specification. *Dev Cell*, 46: 285–301.e9.

Sanchez-Mut JV, Heyn H, Silva BA, Dixsaut L, Garcia-Esparcia P, Vidal E, Sayols S, Glauser L, Monteagudo-Sánchez A, Perez-Tur J, Ferrer I, Monk D, Schneider B, Esteller M and Gräff J. (2018). PM20D1 is a quantitative trait locus associated with Alzheimer's disease. *Nat Med*, 4: 598–603.

Schäfer A, Mekker B, Mallick M, Vastolo V, Karaulanov E, Sebastian D, von der Lippen C, Epe B, Downes DJ, Scholz C and Niehrs C. (2018). Impaired DNA demethylation of C/EBP sites causes premature aging. *Genes Dev*, 32: 742–762.

Seiwert D, Witt H, Ritz S, Janshoff A and Paulsen H. (2018). The nonbilayer lipid MGDG and the major light-harvesting complex (LHCII) promote membrane stacking in supported lipid bilayers. *Biochemistry*, 57: 2278–2288.

Sutandy FXR*, Ebersberger S*, Huang L*, Busch A, Bach M, Kang HS, Fallmann J, Maticzka D, Backofen R, Stadler PF, Zarnack K, Sattler M, Legewie S* and König# J. (2018). *In vitro* iCLIP-based modeling uncovers how the splicing factor U2AF2 relies on regulation by cofactors. *Genome Res*, 28: 699–713.

Tristán-Flores FE, Guzmán P, Ortega-Kermedy MS, Cruz-Torres G, de la Rocha C, Silva-Martínez GA, Rodríguez-Ríos D, Alvarado-Caudillo Y, Barbosa-Sabanero G, Sayols S, Lund G and Zaina S. (2018). Liver X receptor-binding DNA motif associated with atherosclerosis-specific DNA methylation profiles of *Alu* elements and neighboring CpG islands. *J Am Heart Assoc*, 7: e007686.

Uhrig S and Klein H. (2018) PingPongPro: A tool for the detection of piRNA-mediated transposon silencing in small RNA-Seq data. *Bioinformatics*: bty578.

Winzi M, Casas Vila N, Paszkowski-Rogacz M, Ding L, Noack S, Theis M, Butter F and Buchholz F. (2018). The long noncoding RNA *lncR492* inhibits neural differentiation of murine embryonic stem cells. *PLOS One*, 13: e0191682.

* indicates joint contribution # indicates joint correspondence

SCIENTIFIC ADVISORY BOARD



IMB is grateful to the following outstanding scientists for the insight, guidance and advice that they have provided in order to help us become a leading research centre.

GENEVIÈVE ALMOUZZI

Institut Curie, PSL University, Sorbonne University, UPMC University, CNRS, UMR3664, Paris, France



PETER BECKER (CHAIR)

Biomedical Center Munich, Ludwig Maximilian University (LMU), Munich, Germany



BRADLEY CAIRNS

Huntsman Cancer Institute, University of Utah, Salt Lake City, USA



RUDOLF JAENISCH

The Whitehead Institute for Biomedical Research, Cambridge, USA



JOSEF JIRICNY

Institute of Molecular Cancer Research, University of Zurich, Switzerland



RUTH LEHMANN

Skirball Institute, NYU School of Medicine, New York, USA



JAN-MICHAEL PETERS

Research Institute of Molecular Pathology (IMP), Vienna, Austria



RESEARCH ENVIRONMENT

IMB is embedded in a strong and dynamic research environment. It is located on the leafy campus of Johannes Gutenberg University, just west of Mainz city centre.

With 10 departments, more than 150 institutes and 32,000 students, Johannes Gutenberg University is one of the largest German universities. In biomedical research, the university has built strong, interdisciplinary centres dedicated to neuroscience, cardiovascular medicine, immunology and oncology.

The University Medical Centre, which is located near the main university campus, has a strong focus on clinical and translational research and has researchers who also work in close contact with IMB. In addition to the University, IMB has two Max Planck Institutes (the Max Planck Institute for Chemistry and the Max Planck Institute for Polymer Research) and Mainz's University of Applied Sciences as immediate neighbours.

Mainz is also surrounded by a number of towns and cities with extensive research activities. For instance, Frankfurt is only 35 km away and is home to Goethe University, which has over 46,000 students and 10 research institutes within the Biochemistry, Chemistry and Pharmacy Department alone. Furthermore, there are several Max Planck Institutes in Frankfurt (including the Max Planck Institute for Biophysics, the Max Planck Institute for Brain Research and the Ernst Strungmann

Institute for Cognitive Brain Research). In addition to Frankfurt, nearby Darmstadt is home to both a Technical University, whose Department of Biology has a focus on synthetic biology and the biology of stress responses, and a University of Applied Sciences that includes a focus on biotechnology.

Furthermore, there is an extensive industry R&D presence, with, for example, the headquarters of Boehringer Ingelheim and the Merck Group both in close vicinity.



WHERE WE ARE

IMB is located in the city of Mainz, a charming, open-minded city that dates back 2,000 years to Roman times and still has a historic centre with a magnificent medieval cathedral. It was also here, in 1450, that Johannes Gutenberg invented modern book printing. The city is located at the confluence

of two of the most important rivers in Germany, the Rhine and the Main, and has spectacular esplanades. Mainz is within easy reach of both cosmopolitan Frankfurt, with its famous opera house, avant-garde museums and glass-and-steel banking district, and the Rhine valley region with its

castles, vineyards and nature reserves that offer great outdoor activities. With Frankfurt airport – one of the largest airports in Europe – only 25 minutes away, countless European and overseas destinations are within easy reach.




 *iml*
Institut für
Molekulare Biologie

Ackermannweg 4


CAMPUS MAP

OF JOHANNES GUTENBERG UNIVERSITY MAINZ





JOHANNES GUTENBERG
UNIVERSITÄT MAINZ



CONTACT

Institute of Molecular Biology gGmbH
supported by the Boehringer Ingelheim Foundation

Ackermannweg 4
55128 Mainz, Germany

Phone: +49-6131-39-21501
Fax: +49-6131-39-21521

www.imb.de
info@imb.de



IMPRINT

Published by Institute of Molecular Biology gGmbH

Executive editors Helle Ulrich and Ralf Dahm

Coordination Jonathan Byrne and Christine Omeni

Editors Jonathan Byrne and Christine Omeni

Layout and design cala media

Scientific Figures Courtesy of the respective Research Groups and Core Facilities

Staff and funding data Courtesy of IMB Human Resources and Grants Office

Image Credits

Cover: False colour image of a 2 mm section of a chemically cleared mouse brain. Image composed of 35 single images. Sample prepared by Oriane Blanqui (group of Heiko Luhmann, University Medicine Mainz) and imaged by IMB's Microscopy Core Facility. Image credit: Sandra Ritz.

Portraits of Group Leaders and Core Facility heads (p2, 5-40, 44-51), pictures of IMB researchers (p 52, 56-58, 63, 64, 69), IMB building (inside cover, p3, p73, p76-77): Thomas Hartmann

Microscopy images: (p3) Stage 38 *Xenopus* larvae stained for 5hmc (red) and yoyo1 (green). Image credit Victoria Hatch (Niehrs Group). (p4) Stained section of mouse intestine. Image credit Bernadette Mekker. (p61) Monolayer section of whole murine colon after DNA damage. Image credit Konrad Gronke (Diefenbach group, Charite, Berlin), image taken at the AF7000 Fluorescence Microscope (Leica) at 10x magnification by the Microscopy and Histology CF, IMB.

Close-up of magazine pages (p67): G10ck@Shutterstock.com

Journals (p70): Carlos_Castilla@Shutterstock.com


Aerial view of Mainz over the river (p72): Meinzahn@iStock.com

Frankfurt by night (p72): Jotily@iStock.com

Johannes Gutenberg statue (p72): Landeshauptstadt Mainz

SAB (p73): All images courtesy of SAB members

All other images: IMB archive



ANNUAL REPORT 2018
www.imb.de